

PHILADELPHIA MEDICAL TIMES.

A BI-WEEKLY JOURNAL OF
MEDICAL AND SURGICAL SCIENCE.

EDITED BY HORATIO C. WOOD, M.D.

SEPTEMBER 14, 1878.

CONTENTS.

ORIGINAL COMMUNICATIONS:

- TWO CASES OF OPIUM-POISONING—FAL-
TURE OF ATROPIA TREATMENT. By
FRANCIS L. HAYNES, M.D. . . . 577
- SUCCESSFUL CASE OF TREPHINING FOR
EPILEPSY. By Prof. D. HAYES AGNEW . . . 578
- UNGUENTUM HYDRARGYRI OXIDI FLAVI.
By J. T. WALKER, M.D., Ph.D. . . . 580
- CHLORAL HYDRATE AND OXIDE OF ZINC
IN ACUTE INTESTINAL DISEASES OF
CHILDHOOD. By JAMES L. TYSON, M.D. . . . 581
- A SINGULAR CASE OF INFANTILE CON-
VULSIONS. By J. F. WALSH, M.D. . . . 582

NOTES OF HOSPITAL PRACTICE:

- COLLEGE OF PHYSICIANS AND SUR-
GEONS, NEW YORK—Clinic of Prof. A.
JACOB, for Diseases of Children: Hereditary
Syphilis, with Osteitis and Pulmonary Infiltration
—Prolapse of the Rectum . . . 582

TRANSLATIONS:

- Bloodless Operations upon the Tongue—Symptoms
and Treatment of Psoriasis Universalis—Injection
of Tincture of Iodine into the Knee-Joint—Uterine
Myoma; Subacute Peritonitis following Simple Vag-
inal Examination; Death—Penetrating Wound of
Abdomen; Recovery—Prevention of Hemorrhage
after the Use of Esmarch's Bandage . . . 584, 585

EDITORIAL:

- MARINE HOSPITAL SERVICE . . . 586

CORRESPONDENCE:

- LONDON LETTER . . . 587
- A QUESTION OF PRIORITY . . . 591

PROCEEDINGS OF SOCIETIES:

- PATHOLOGICAL SOCIETY OF PHILADEL-
PHIA:* Cancer of the Right Kidney; Thrombosis
of Ascending Cava and Left Renal Vein—Colloid
Cancer of the Abdomen—Case of Recurrent Fibroid
of Axilla—Three Cases of Heart Disease associated
with Kidney Disease—Cancer of the Ascending
Colon—Embolie Pneumonia . . . 592

REVIEWS AND BOOK NOTICES:

- Handbook of Ophthalmology. By Prof. C. Schweig-
ger, of the University of Berlin. Translated by
Porter Farley, M.D. . . . 598
- Diseases of the Nervous System. By Samuel Wilks,
M.D., F.R.S. . . . 599

GLEANINGS FROM EXCHANGES:

- Unusual Occurrence in the Attempt to reduce a Dis-
located Humerus—Opium Habituation—The Ac-
tion of Sulphuretted Waters in reproducing the
Symptoms of Syphilis when Latent . . . 599

ARMY NEWS:

- Official Changes from August 25 to September 7, 1878 600

TERMS.—Yearly Subscription, \$4.00, in advance. Single Number, 20 Cents.

J. B. LIPPINCOTT & CO., Publishers,
715 and 717 Market St., Philadelphia.

LONDON: 16 SOUTHAMPTON ST., COVENT GARDEN.

Special Agents:

{ E. R. PELTON, 25 Bond St., New York.
A. ROMAN & CO., San Francisco.

A. WILLIAMS & CO., Boston.
BOSSANGE & CIE., Paris.

EXTRACT OF MALT

WITH COD LIVER OIL.

Our Improved Extract of Malt with 50 per cent. of our fine Norwegian Oil is pre-eminently **THE** emulsion of Cod Liver Oil, since it *has double the nutritive effect of any other emulsion of oil*. It not only contains the largest amount of nutritive food in the smallest bulk of liquid, but by the action of the Diastase contained in the malt, it is capable of transforming many times its bulk of amylaceous or graminaceous food into saccharine, thus rendering it easily digestible even by the most delicate stomach of the invalid. The disagreeable eructations from the stomach, which constitute one of the greatest objections to the use of all preparations of Cod Liver Oil, are completely obviated in this preparation. We would direct the attention of physicians to the special fact that it **UNITES PERFECTLY WITH WATER IN ANY PROPORTION**, and that the oil does not again separate, thus showing how perfectly the oil is emulsified.

DOSE.—A Dessert to Tablespoonful immediately after Meals.

Prepared with the most scrupulous care by

KEASBEY & MATTISON,

Manufacturing Chemists,

332 N. Front Street, Philadelphia.

SEND FOR PAMPHLET ON MALT AND NEW REMEDIES.

Soluble Medicated Intra-Uterine Pencils,

FOR THE TREATMENT OF CHRONIC DISEASES OF THE UTERUS.

These pencils being soluble at the temperature of the body, gradually melt when introduced into the womb, and thus bring the medicament into thorough and prolonged contact with the diseased surface.

THEY ARE MEDICATED WITH

Ergotin, 5 and 8 grains.

Nitrate Silver, 5 grains.

Sulphate Zinc, 1, 2, 4, and 6 grains.

Iodoform, 2, 3, and 5 grains.

Carbolic Acid, 1, 2, and 4 grains.

Ext. Opium Aq., 1 and 2 grains, etc.

Price per box of half dozen

\$0.50

Price per dozen boxes

\$4.50

CHARLES L. MITCHELL, Pharmacist, Ninth and Race Sts., Philadelphia.

SENT BY MAIL TO ANY ADDRESS.

Send stamp for circulars and samples.

OUR MAKE OF SUGAR-COATED PILLS

Received the Highest Award attainable,

A Centennial Medal and Diploma,

For Superiority of Finish and Purity of
Ingredients.

In prescribing, please specify (B. & C.)

**BULLOCK & CRENSHAW,
PHILADELPHIA.**

VINUM DIGESTIVUM. DIGESTIVE WINE.

A vinous combination of Pepsin, manufactured by a recently improved process, furnishing the most efficient and powerful liquid preparation of this invaluable remedial agent yet offered.

One fluidounce will dissolve at least 500 grains of coagulated albumen.

Containing the digestive principle of the animal stomach in a form adapted for immediate effect, it commends itself as the remedy for Dyspepsia, Indigestion, and the different phases of weak or disordered stomach.

DOSE.—One or two teaspoonfuls immediately after eating.

ORIGINATED AND MANUFACTURED BY

WM. PROCTER, JR., CO.,
S. W. Cor. Ninth and Lombard Sts.,
PHILADELPHIA.

BLAIR'S SPECIALTIES.

BLAIR'S

PREPARED WHEAT FOOD.

A Nutritious Diet for Infants and Invalids.

Contains a larger proportion of Phosphates and Nitrates (muscle- and bone-forming elements) than most farinaceous foods. Sample packets free. Try it.

BLAIR'S

MEDICATED PRUNES.

Cathartic and Laxative.

Very pleasant to the taste, and certain in effect, without griping. Dose, one to three, as required.

RUBBER FELT.

This is not the "impervious lint," consisting of lint with a sheeting of rubber glued on, but has been prepared expressly for us, the same as rubber cloth. Very superior for the application of poultices, liniments, etc., where it is desired to keep in the moisture and keep the part warm.

THE EUREKA INHALER.

A convenient apparatus for the application of muriate of ammonia (fumes) and volatile remedies to the nasal cavities, throat, and lungs. The advantages of our inhaler are low price, simplicity, durability, and compactness.

SEND FOR CIRCULARS.

MADE BY

HENRY C. BLAIR'S SONS,
APOTHECARIES,
Walnut and Eighth Sts., Philadelphia.

PHILADELPHIA MEDICAL TIMES.

PHILADELPHIA, SEPTEMBER 14, 1878.

ORIGINAL COMMUNICATIONS.

TWO CASES OF OPIUM-POISONING —FAILURE OF ATROPIA TREATMENT.

BY FRANCIS L. HAYNES, M.D.,

Assistant-Surgeon to the Episcopal Hospital.

I.—HALF AN OUNCE OF LAUDANUM SWALLOWED AND RETAINED—RECOVERY.

SAMUEL S., aged 50, a weaver, was a man of eccentric character and intemperate habits. He had been for years afflicted with phthisis, and was greatly emaciated. On August 9, 1878, during a debauch, which had commenced five days previous, he swallowed half an ounce of laudanum. Vomiting did not occur at any time. The preparation was not the official one, but I was assured by the very careful apothecary from whom it was purchased that it contained the active principle of nine grains of opium (to the half-ounce of menstruum). The patient was taken to a police-station, and was treated by electricity and forced exercise. I did not see him until 4½ hours had elapsed.

4½ hours.—He was very sleepy, but could be awakened by shaking; he answered questions, and could walk, with the assistance of two men. He would not swallow, or allow the stomach-pump to be used. Respiration 6, somewhat stertorous; pulse 156, weak; pupils—right one-half dilated, left extremely contracted; both immobile. Atropia sulph. gr. $\frac{1}{15}$ was injected under the skin. He was aroused at intervals by shouting, whipping, and shaking, and by forcing him to walk, but did not breathe more frequently when awake. The stertor disappeared when he was awake.

5 hours.—Condition the same. Atropia sulph. gr. $\frac{1}{15}$ injected.

5½ hours.—He could scarcely be awakened. When attempts were made to exercise him, his legs "doubled up." Respiration 5, stertorous; pulse 158, weaker.

5¾ hours.—Atropia sulph. gr. $\frac{1}{38}$ injected.

6 hours.—He could not be aroused. The left pupil was somewhat larger; the right was of the same size as at 4½ hours. Respiration and pulse same as at last note. Atropia sulph. gr. $\frac{1}{38}$ injected.

6¼ hours.—Respiration 5, stertorous; pulse 160, very weak. Left pupil still larger; right continued at same diameter.

6¾ hours.—He was completely comatose. Respiration, pulse, and pupils the same. Atropia sulph. gr. $\frac{1}{38}$ injected.

8 hours.—He was apparently dying. The lips were blue, the surface cold and clammy, the eyes turned up. Respiration 4, very noisy; pulse almost imperceptible, and so frequent that it could not be counted. The left pupil

had dilated to the same extent as the right, which retained its original diameter. He remained in this condition for half an hour, then gradually improved.

9½ hours.—He could be readily aroused. Respiration 5, less stertorous. Pupils and pulse same as when last noted.

15½ hours.—He was awake and troubled by hallucinations, but was occasionally sensible. Respiration 11, quiet; pupils same; pulse 144.

24 hours.—Delirium since last note; sleeplessness; dryness of throat. Respiration 20; pulse 140.

43 hours.—He was still sleepless, and was troubled at times by hallucinations. A sleeping-draught was ordered.

From this date the history of the case is one of pneumonia occurring in a consumptive. He succumbed August 18.

The total quantity of atropia sulph. given was gr. $\frac{4}{15}$.

Remarks.—Atropia was not given in this case as an antagonistic poison, but as a stimulant to the respiratory centres. During its use all the symptoms were aggravated; and one and three-fourths hours elapsed, after the last dose, before any improvement occurred.

Atropia is supposed to be beneficial in opium-poisoning—(1) by antagonizing in some mysterious and undefined way the action of opium, just as quinia antagonizes the action of malaria on the system;* (2) by causing the dilated and paralyzed vessels of the brain and spinal cord to contract, and thus obviating death by congestion or œdema of these organs;† (3) by stimulating the heart, and preventing death by cardiac exhaustion;‡ (4) by stimulating the respiration, and thus combating the most frequent cause of death in narcotic poisoning,—failure of respiration.§

In this case the atropia had an opportunity to act in any of these ways. The dose was large enough to act as an antagonistic poison,|| and not too large to act as a cerebral vaso-motor, a cardiac, or a respiratory, stimulant.

For the history of the following case I am indebted to Dr. Todd, of West Point, Ga. It has not, I think, been published hitherto.

* Norris, Am. Journ. Med. Sci., Oct. 1862; Johnston, Med. Times and Gaz., Sept. 7, 1879, p. 268.

† Bennett, Antagonism of Medicines, Lond., 1875, p. 46.

‡ Harley, Old Vegetable Neurotics, p. 309.

§ H. C. Wood, Am. Journ. Med. Sci., April, 1873, p. 339.

|| Nine grains of opium, the dose taken in this case, are about equal to one grain of morphia, a dose which is sometimes used therapeutically, by the mouth, or injected under the skin. According to the ordinary doses employed, gr. 4-19 of atropia should at least equal as a poison the above-mentioned dose of morphia.

II.—ONE OUNCE OF OPIUM TAKEN—A PORTION
VOMITED—DEATH.

B. R., aged 56, a merchant, had been at intervals for eighteen months a sufferer from attacks of insanity, during which he had attempted by various means to commit suicide. On December 20, 1872, he ate one ounce of gum-opium.

1 hour afterwards.—He was slightly sleepy. Refused treatment.

2 hours.—The drowsiness was increasing. Vomiting of a quantity of opium followed the administration of sulphate of zinc, through the nose, by means of a funnel and catheter. Forced exercise was used. Hypodermic injections of veratrum viride tincture, eleven drops in a drachm of whisky, were commenced, and continued at intervals of half an hour.

7 hours.—Hitherto the patient had not been very drowsy. He had given rational answers to questions; the respiration had not fallen below 14, or the pulse below 65, a minute. The pupils had not contracted. "At this time Dr. X., the family physician, arrived, and only one more dose of veratrum viride was given. The patient recognized Dr. X. Dr. X. counted his pulse and respiration, which were as given above. We commenced to use the cold douche.

"7½ hours after the ingestion of the poison we commenced to inject atropia under the skin, in doses of gr. $\frac{1}{20}$. This was repeated every fifteen minutes. From the first dose the coma deepened and the respiration became less frequent.

"8 hours and 20 minutes.—The pupils began to dilate.

"8½ hours.—At this moment, precisely, he ceased suddenly to breathe. The heart continued to beat 65 a minute for several minutes after the breathing had ceased. Artificial respiration and electricity were tried, but in vain."

Remarks.—Atropia certainly did not stimulate the respiration in this case, the patient dying from failure of that function. (It is not certain, from the notes, whether four or five doses were injected; but probably only four, as respiration ceased just as the moment came for the fifth injection. The state of the pupils, though this is not a very important point, is not very satisfactorily indicated.)*

* The subject is one of such importance that the editor of the *Times* takes the liberty of pointing out that, according to his reading, these cases do not disprove the value of atropia in opium-poisoning. In the last case the patient took the equivalent of nearly thirteen ounces of laudanum: of course in immediate evacuation of the stomach lay the only chance of escape. In the first case, apparently, not enough of atropia was used to so far counteract the effect of the laudanum as to produce any immediate effect. Only after the lapse of some hours does the report afford any indication of belladonna symptoms. There can be no definite relative dose of the two poisons, as different individuals offer very different susceptibility to the two narcotics.

SUCCESSFUL CASE OF TREPHIN-
ING FOR EPILEPSY.

BY PROF. D. HAYES AGNEW.

(Reported by W. HOBSON HEATH, M.D., Ex-Resident Surgeon University Hospital, Resident Physician Philadelphia Hospital.)

M. R. B. S., æt. 45 years, was wounded in Tennessee on January 1, 1863, by a piece of shell which exploded about twenty feet above him, a fragment striking him upon the top of the head and chipping off a piece of skull. He lost consciousness for about an hour, was taken to a hospital and placed with other wounded. No evidence of compression. The injury was not followed by any inflammation of the brain or meninges, although the wound was attacked by erysipelas two weeks after. He rejoined the Confederate army eight months after receipt of the injury, during which time several fragments of bone were discharged; his health otherwise being perfect. The first attack of epilepsy came on suddenly before breakfast, six years later (1869), upon a damp, inclement morning. There was no distinct aura; the fit was violent, and the following stupor deep, and it was the most violent of all his seizures. The cicatrix had never given any neuralgic trouble or become inflamed.

Two months later a second seizure occurred of the same character, and very severe, and up to the present time they have recurred at periods varying from one to six weeks. At one time they were held in check by large doses of the "bromides." The past year the attacks have been preceded by an aura, which commences on the inner and under side of the arm near the wrist, and sometimes at the tips of the little and ring fingers, and is described as a sleepy, numb feeling. The disease has seriously interfered with his business, impaired his mind and memory, especially during the past two years, and on several occasions jeopardized his life.

General health good, habits excellent, is married, and children healthy.

Parents healthy, and habits good. No history of epilepsy in family.

Examination of head shows a cicatrix two and three-quarter inches long over right parietal bone. The cicatrix is depressed one and three-quarter inches of its length at one end. Depression measures one and a half inches in length at the bottom, one-third of an inch wide, one-third of an inch deep.

This patient had been in the hands of many of the best physicians in the South, without relief. He applied to Prof. Agnew for surgical interference to remedy his condition. He was kept quiet for a few days, and his general condition was looked after; had no fits for thirteen days prior to admis-

sion. At the time of operation he was etherized, his head was shaved, and a crucial incision made over the cicatrix, each flap being dissected off from the pericranium. The somewhat free hemorrhage was controlled by the flaps being held and the vessels compressed by the assistants. The trephine, an ordinary fluted one, the size of a twenty-cent piece, was applied over and included one end of the depression, and was very slowly worked. It was some twenty minutes before the skull was sawed through, and then upon examination it was found to have been perforated only at one-half of the circumference of the trephine. This irregularity was due to the fact that the bone, especially the inner table, had become hypertrophied at and around the point of injury, and the point first sawed through was where the process had been less—farthest off from the injured portion of bone.

Examination of the pieces removed and of the skull itself showed that there was no depressed fracture at all; that the injury sustained by the bone was simply the chipping off of a piece from the outer table; that, in consequence of this, the bone, particularly the inner table, had become hypertrophied, the thickest part being apparently around the injury, and therefore the pressure upon the brain was general over a surface probably many inches around. Two discs of bone, which included the *depression in the bone*, were removed, and showed the condition stated, and also that the increased thickness was of such extent that it was not thought prudent to remove it entirely. The fact that frequently in trephining the instrument passes through to the dura mater at one point before it does at another (and it was so especially in this case, though from pathological reasons) suggested to Prof. Agnew the idea of having a trephine serrated only on a portion of its circumference, the remainder being smooth. Such an instrument could be used without injury to the membranes where they were exposed, and in the present case would have been very useful. On removing the pieces of bone the membranes were found uninjured, and there was no trace of previous inflammation. The flaps were closed over the wound, but no stitches or adhesive plaster applied to keep them in place, and thereby exert any pressure or obstruct the drainage. The treatment then ordered was a plain water-dressing locally, and calomel $\frac{1}{2}$ grain, opium

$\frac{1}{6}$ grain, every two hours, with an occasional dose of bromide of potassium, if restless, milk diet, elevation of the head, and perfect quiet.

Patient came out of ether nicely, and did not vomit; complained slightly of headache; had no chill, no increase of temperature, and slept well under bromide of potassium until the next morning, when he was taken with a convulsion, not very severe or long, which was attributed to the operation.

The case after this did remarkably well. There was no inflammation of the brain or meninges, though it was expected to follow.

The temperature during the first week arose to 100° twice; after that it remained at 98° , 99° , until the patient was discharged. The pulse, never above 100, and averaging 90, was throughout soft and regular. The tongue was slightly furred, due to the irregularity of the bowels and confinement. The constipation was obstinate, owing to the recumbent position and a natural torpid habit. Seidlitz powders and the occasional judicious administration of an enema only were used.

The appetite was good after the first four days.

The most annoying symptom complained of throughout was headache. This was constant and frontal,—right over the eyes; it diminished after the first week, and assumed an intermittent character, being produced, apparently, or influenced most by constipation. At one time it appeared like neuralgia of the supra-orbital nerve, and aconitia ointment was applied, and gave relief.

Bromide was given throughout the treatment; the calomel and opium were discontinued after the first week. The wound was dressed with plain water-dressing, and nothing was allowed to interfere with free drainage. After the granulation reached the level of the scalp, the edges of the flaps were gently approximated by strapping.

At the end of twenty days he was allowed to sit up in bed, and by the thirty-second day to get out of bed, remaining up longer day by day.

The operation was not followed by any exfoliation of bone.

The patient was discharged six weeks after operation, to return to the South.

This makes the second case of trephining for epilepsy by Prof. Agnew at the University Hospital, within two years, with

the recovery of both patients and the disappearance of the malady.

The most peculiar and interesting feature in this case was the general thickening of the bone around the seat of injury and the relief by the operation, though the entire amount thickened and pressing upon the brain was not removed.

The rapid and uninterrupted convalescence is worthy of note, considering his age and condition, and the gravity of the operation.

UNGUENTUM HYDRARGYRI OXIDI FLAVI.

BY J. T. WALKER, M.D., PH.D.

IN the *Medical Times* of July 20, 1878, there appeared an article written by Dr. Landesberg condemning the above ointment, for its non-stability and difficulty of preparation, in the following language: "1st, it is *very liable* to decomposition; 2d, it is very hard to *triturate*. Whatever constituents may be used for the ointment, decomposition sets in sooner or later. The ointment, which, if properly attended to, is of a fine yellow color, becomes, after a short time, dirty yellowish, smells rancid, and loses its healing power."

He says further, "How difficult it is to *triturate* the yellow oxide of mercury is proved by the fact that the correct composition of the ointment is found with only few druggists," etc.

How the doctor could arrive at the above conclusions is difficult for one to understand, unless he has repeatedly fallen into the hands of ignorant or unprincipled druggists.

In the first place, the ointment is not very liable to decomposition if made with suitable material; secondly, it requires no *trituration*, as the yellow oxide is an amorphous powder, prepared by precipitation, showing no crystalline structure even under the microscope; in this respect totally unlike the *red* oxide.

On the 24th ult. (more than four weeks ago), Mr. D. H. Latham, Jr., prepared for me the following ointments (all of which are in a good state of preservation), taking no more than ordinary care in preparing them, and taking special care to give them the usual exposure by removing the lids from the jars daily and allowing access of

air. They were dispensed in an ordinary half-ounce glass jar with a wooden lid, the same as used by most apothecaries in dispensing ointments.

No. 1.

R Hydrargyri oxidi flavi, gr. xxx;
Unguenti benzoini (U.S.P.), ʒiiss.

No. 2.

R Hydrargyri oxidi flavi, gr. xxx;
Unguenti petrolei (Cosmoline), ʒiiss.

No. 3.

R Hydrargyri oxidi flavi, gr. xxx;
Unguenti aquæ rosæ, ʒiiss.

No. 4.

R Hydrargyri oxidi flavi, gr. xxx;
Unguenti aquæ rosæ, ʒiiss;
Balsam. Peruv., gtt. iii.

No. 5.

R Hydrargyri oxidi flavi, gr. xxx;
Unguenti cum oleo ricino, ʒiiss.

It will be observed that the amount of yellow oxide used is greater than ordinarily prescribed, which would be more favorable than otherwise for oxidation. No decomposition has yet taken place in any of the ointments, but of course would in time, with the exception probably of that made with *cosmoline*, it being no doubt by far the best material for all ointments, and should be preferred where an ointment would be required to be kept a very long time; but a month is quite as long, even longer, than any physician expects a patient to use an ointment without renewal.

No. 3 was made with cold cream (rose-water omitted). This cold cream was not made for the oxide of mercury ointment, but was taken from some that had been made two weeks, the object being to use no more than *ordinary* care in preparing and selecting material, so as to come to an impartial conclusion concerning the above ointments.

No. 4 was made with same cold cream (U. S. P. formula), minus rose-water.

No. 5 was made from a recipe furnished by Mr. Henry A. Bower (*vide Journal of Pharmacy*, Phila., vol. xlii. p. 303). An ointment of the red oxide of mercury, prepared thus, I have seen keep for six months without undergoing any change. This probably is almost as permanent as *cosmoline*. The following is the formula:

R Hydrargyri oxidi rubri (vel flavi), 5x;
Olei ricini, fʒi;
Adipis (lat.), ʒvii;
Cera flavæ opt., ʒii.

Melt the wax and lard together, and mix

with the castor oil. On cooling, add the precipitate in very fine powder, stir constantly with a wooden spatula until cold. From the above experiments I am convinced that no one is worthy of the title apothecary who cannot furnish a satisfactory ointment of yellow oxide of mercury.

201 COLUMBIA AVENUE, PHILA.

CHLORAL HYDRATE AND OXIDE OF ZINC IN ACUTE INTESTINAL DISEASES OF CHILDHOOD.

BY JAMES L. TYSON, M.D.

HIGH heat for the past six or eight weeks, together with the irritation of dentition and improper diet, or over-feeding, or both, have been prolific factors in the generation of that troublesome and not rarely fatal malady among children, in ordinary parlance cycled summer complaint, whether it presents itself under the form of simple diarrhea, cholera infantum, entero-colitis, or dysentery. Though the unwholesome atmosphere of a city in hot weather contributes largely to its production and fatal result, I am inclined to think that unsuitable food and too much of it may in the country, from my observation of both localities, be pronounced almost as frequent a cause of its prevalence and fatality there as are the foul airs in the crowded alleys of a populous town. At all events, it has prevailed to a considerable extent in this county, and, as I have had a good many cases under my care, some of which were almost *in extremis* when first seen, I have been urged to present a brief abstract of the treatment instituted, though by no means new to many, which in my hands has resulted so satisfactorily. I am more particularly induced to refer to it now, from having read a note addressed to the editor of the *Medical Times*, by Dr. W. L. Newell, of Millville, New Jersey, announcing the benefits which resulted from chloral enemata in his and his colleague's hands in cases of dysentery in adults. It is the employment of this agent in that form, along with other treatment, among children within the year,—say from six to nine months old,—whose claims to consideration I advocate and desire to enforce with all the earnestness that its merits demand. I was much gratified to learn from Dr. Newell's note that his treatment for adults so fully vindicated

my preconceived impressions of its utility in cases of children. Simple attacks of the complaint in this vicinity readily yielded to a change or diminution of diet, and a cold bath two or three times a day. Others were of a much graver type, the discharges being lenteric, mixed with blood or bloody water, from twelve to twenty occurring in the course of a day, and in some cases the tenesmus so excessive that the moment an enema was administered it was expelled. This spasmodic action of the sphincter and lower bowels could only be controlled by repeated resort to the remedy, two or three applications being requisite before it could be retained, and then only by directing the nurse or mother to compress the glutæi muscles on either side, close over the anal orifice, for two or three minutes. When thus kept in immediate contact with the inflamed, sensitive, and irritable tissue, the benefits were prompt and enduring. Tenesmus, or choreal spasm of the bowel, was arrested, pain and inflammation were allayed, and the little sufferer would rest or sleep comfortably for several hours. A repetition of the enema was made once, sometimes twice, in the twenty-four hours, with increased comfort and alleviation of all the symptoms.

In cases of this kind, as well as others, of course great attention was paid to the preparation of food, so that entozoa, infusorial, or bacterial spores in the fluid used were thoroughly *sterilized*. To accomplish this the milk was added to *boiling water* containing a little gelatin and arrow-root. The milk should not be boiled. A teaspoonful of lime-water should be put in every teacupful of this preparation, which should always be given cold. The amount of chloral used in each application was about two grains dissolved in one or two teaspoonfuls of starch water.

Along with this local treatment, two grains of oxide of zinc and three of lactopeptine in mucilage were given every five or six hours. This combination exerts a happy influence on the primæ viæ, enabling the child to digest its food more thoroughly, and controlling the number while it alters the character of the evacuations in a day or two. We are indebted to Dr. Brackenridge, of Edinburgh, for having first suggested the use of oxide of zinc in these maladies of children,—a detailed statement of cases in which he had successfully employed it being pub-

lished,—and to a more recent article in the *Glasgow Medical Journal* by Dr. I. Crawford Renton and inserted in the last number of *Braithwaite*. Some of the cases in which the foregoing treatment was carried out were of a desperate, apparently hopeless, character, but in all benefit was soon apparent, and the little patients recovered. In no instance was any preparation of opium or calomel resorted to; but I can well understand the advantages claimed for minute doses of mercuric chloride in a well-written article on “Acute Intestinal Catarrh of Infants,” from the pen of Dr. Rudolph Ravenburg, of Washington, D.C., published in a recent number of a New York medical journal; and many cases might occur in which its employment, alone or in conjunction with the treatment already detailed, might be clearly indicated.

Subjoined are the formulas used in the cases referred to.

I may add that the cold bath three times a day was invariably insisted upon, but never at a lower temperature than 80° to 85°.

R Chloral hydrate, ʒss;

Starch water (amylum), ʒij.

M. Ft. solut.

Sig.—Enema. One to one and a half teaspoonfuls to be *thrown* into the bowel from a small glass syringe.

R Zinci oxidi, ʒss;

Pulv. g. acaciæ et sacch. alb., āā ʒij;

Lactopeptin, ʒj;

Aq. cinnam., q. s. ut ft. ʒi.

M. S. A.

Sig.—A teaspoonful every five or six hours.

MONTGOMERY COUNTY, August 16, 1878.

A SINGULAR CASE OF INFANTILE CONVULSIONS.

BY J. F. WALSH, M.D.

AUGUST 22, 1878, I was called in great haste to see M. C., æt. 6 years, when I found the boy in violent convulsions. He was apparently a healthy lad, and on inquiring into his previous history I learned that he had never been subject to fits, nor had he ever had any serious sickness. He had never been troubled with worms.

His mother said that he had been found, about two hours previous, in an open lot in front of a grocery-store, in an insensible condition. Strewed around him was a quantity of spoiled raisins which the grocer had thrown out. He was taken into the store, where he

vomited a few raisins. About a half an hour later, 11½ A.M., he was removed home. My office-hours not commencing until 1 P.M., the family did not send for me until that time. In the mean while he had been in a state of almost continuous eclampsia, having but few remissions. He had not been given any medicine before my arrival; all the mother did was to put him in a warm bath, which, however, did not relieve the symptoms.

When I got to the house his condition was as follows. Whole body convulsed. Unconscious. Cheeks dusky red,—face bathed in sweat. Pupils dilated and insensitive to light; strabismus. Breathing labored. Heart acting very irregularly, fast (one hundred and fifty per minute), and feebly. Slight frothing at corners of mouth. Grinding of teeth. Belly tympanitic.

Believing that the raisins he had eaten were the source of the trouble, I administered, about fifteen minutes after my arrival, during a remission, three teaspoonfuls of wine of ipecac. About a half an hour later he was seized with violent vomitings, and threw up a large quantity of partially-chewed raisins. Immediately the convulsions ceased, he went off into a quiet sleep, and subsequently awoke quite well.

CAMDEN, N.J.

NOTES OF HOSPITAL PRACTICE.

COLLEGE OF PHYSICIANS AND SURGEONS, NEW YORK.

CLINIC OF PROF. A. JACOBI, FOR DISEASES OF CHILDREN.

(Reported by P. BRYNBERG PORTER, M.D.)

Concluded from p. 560.

HEREDITARY SYPHILIS, WITH OSTEITIS AND PULMONARY INFILTRATION.

ROSANNA G., 9 years of age. Since she was one year old the child has suffered from glandular lesions about the neck and throat, and several times abscesses of this character have been lanced. Up to two years ago she had no nasal catarrh, as far as can be ascertained, but since then she has been very seriously troubled in this manner. There has been an almost constant discharge of pus and blood, with particles of decayed tissue,—and almost from the first, the mother says, the septum of the nose has been affected. The nasal bones have also become implicated, and you notice that there is now considerable deformity resulting. No very clear history of syphilis can be obtained from the mother, as she denies ever having had the ordinary symptoms, but the fact that she has had no less than six miscarriages would leave little ground for doubt that the child now before

us is suffering from hereditary syphilis. One other child besides this one was born at full term, but is now dead. The mother says that it never had any eruption on it. One of the abortions was at the eighth month, and all the others at a considerably earlier period of gestation. Nothing is known in regard to the father.

Perforation of the septum of the nose is almost universally regarded as an evidence of syphilis. Ordinarily, I believe that it is, but at the same time I have seen several instances in which this was not the case. Sometimes the ulceration from which perforation finally results is very superficial at first, but, on account of the peculiar character of the cellular tissue, it goes on slowly increasing in extent and depth. A perforation originating in this manner is usually in the lower part of the septum. Osteitis and periostitis are very common manifestations of hereditary syphilis, and where syphilis has been partially arrested in the parent, bone-disease is frequently the only evidence of the affection seen in the child. Osteitis is more often met with than periostitis in hereditary syphilis, while the latter, as you know, is more commonly met with in the acquired syphilis of the adult.

There is another feature of this case which is quite striking, and that is the unusual prominence of the veins of the neck. This undoubtedly indicates some trouble either with the lungs or the heart, and we will, therefore, proceed to make an examination of the thorax. When the child is stripped you perceive that she is considerably emaciated, and percussion shows that there is marked dullness over a considerable portion of the chest upon both sides. This is more marked over the lower part of the left lung posteriorly and the upper part of the right lung anteriorly. This is certainly not due simply to any pleuritic trouble; for if that were the case there would not be so much obstruction to the circulation as we find present here. On auscultation it is found that there is diminished respiration over a great part of the chest, with some rude breathing at the right apex. When we examine the heart we find that the area of cardiac dulness is much increased, extending farther to the right than normal, and that there is increased shock, while the apex-beat is more to the right than usual. There is, therefore, hypertrophy of the right heart, as we would naturally expect from the condition of the lungs,

which is no doubt chronic; and this is due to the fact that it has more work to do than if the lungs were healthy.

It seems, then, altogether probable that we have to deal here with a specific infiltration of the lungs, and that this has probably existed for years. This condition not infrequently leads to phthisis, and I have not the slightest doubt that a considerable number of cases of phthisis, especially in the cities, originate in the same manner. Hence it is that physicians who are accustomed to meet this class of cases, and who have seen such good results follow a course of mercurial treatment as is frequently the case, have got the idea that mercury is beneficial in all sorts of phthisis. Such, as it seems to me, must be the basis of the opinion that a mercurial treatment is of service in phthisis, and this is, of course, an error; for, while the use of mercury constitutes the best treatment for syphilitic phthisis, it would be about the worst sort of treatment for ordinary tuberculosis.

Our diagnosis in this case, therefore, is syphilitic osteitis and periostitis, with syphilitic pulmonary infiltration; and, under the circumstances, it becomes necessary for us to adopt a persistent but careful course of treatment by mercury. The iodide of mercury produces so much irritation of the alimentary tract as to render its use highly objectionable, and, though the action of calomel is considerably more mild, it is liable to cause too much laxity of the bowels. The bichloride will probably be the best form in which to give the mercury here, and the fortieth or fiftieth of a grain will be about the right dose for the child. Thus, we may order a prescription of one grain of the bichloride to five ounces of water (with a little syrup, if you choose), and of this the child should take a teaspoonful three times a day, about an hour after eating. With this a mouth-wash, composed of half an ounce of chlorate of potassium to the pint of water, should be ordered, and the latter may be employed ten or fifteen times through the day. The gums should be watched carefully, although salivation does not result so frequently in children as in adults who are taking mercury. Still, the drug sometimes produces unpleasant results, and its effect should always be noted with care. If our diagnosis is really correct, and the treatment is faithfully carried out, in the course of two

or three months we ought to expect to find considerable improvement in the condition of the lungs, and, as a result of that, a diminution in the size of the jugular veins.

PROLAPSE OF THE RECTUM.

The last patient whom I shall show you to-day, gentlemen, is a little boy three and a half years old. The mother states that somewhat more than a year and a half ago she noticed that the bowel began to protrude after defecation. The tumor thus formed by it soon became as large as an egg, and is now of even considerably greater size than that. It was formerly of a deep-red color, but is now of a much paler hue. It is a significant point in the history, that the child suffered from diarrhoea during all the summer preceding the prolapsus.

The mother tells me that during this time he often had ten and twelve movements of the bowels in a day, and that the evacuations were copious and passed without straining. The last fact would seem to indicate that there was no local cause for the diarrhoea, like a catarrh of the rectum. In this condition there is an over-secretion of mucus and serum, and sometimes a little blood is passed, whence it is frequently described as catarrhal dysentery.

It is probable that there is no polypus here, but that we have to do simply with a well-marked prolapsus of the rectum. When the bowel thus habitually protrudes after defecation, certain changes take place in its walls. There is a swollen condition of the follicles, and also of the mucous membrane, which becomes much thickened, and there is a rapid throwing off, as well as reproduction, of the epithelium. There is an increased secretion of mucus, as one would naturally suppose, and not infrequently there is some paralysis resulting. When the bowel has come down a number of times, the sphincter ani muscle grows accustomed, as it were, to the protrusion through it, and the bowel becomes swollen all the more after passing it.

In a recent case of prolapsus of the rectum the injection of ice-water is often sufficient to effect a cure, but where this fails we may resort to zinc or alum. One of the best applications is the nitrate of silver, either in stick or in solution of greater or less strength. In whichever form it is employed (and even if only a weak solution is used), it should be immediately neutral-

ized, because otherwise it is almost certain to give rise to extreme tenesmus, and when this occurs it only aggravates the difficulty instead of assisting in its removal. Finally, the actual cautery can sometimes be used with good effect in these cases; and when there is only a paralysis of the sphincter ani, there is nothing of so much service as *nux vomica* or *strychnia*.

Where we have the opportunity (as in private and hospital practice), the most prompt and efficient method of accomplishing a cure, by means of the latter, is to employ it hypodermically, using about the one-sixtieth of a grain every day in this manner.

In this case, when we come to look at the parts to-day, we see that the bowel is now entirely up within the anus. On introducing my finger, after oiling it, into the rectum, I find that there is no polypus present, but that the bowel is everywhere soft and flabby. When the child has a passage, the mother says, the bowel begins to come out before the faeces appear; and this shows that the sphincter is not in a condition to withstand any pressure upon it. When the rectum is down it protrudes fully three inches from the body, she tells me. By way of treatment here I shall order, in the first place, a large injection of cold water every night and morning, which will have the effect of keeping the passages soft, and so less liable to force the bowel down, and will also, I trust, give more tone to the rectal walls. In addition, I will advise the use of an ointment composed of a drachm of the alcoholic extract of *nux vomica* rubbed up with an ounce of fat. Of this the mother is to apply a piece about the size of a bean whenever the bowel comes down, or three times a day whether it protrudes or not.

TRANSLATIONS.

BLOODLESS OPERATIONS UPON THE TONGUE.—Langenbuch (*Chl. f. Chirurgie*, 1878, p. 400; from *Arch. f. Klin. Chir.*) operated in two cases, as follows. The tongue being drawn as far as possible out of the mouth, a long curved needle, armed with a thread, was inserted perpendicularly into the back of the tongue in the median line, some two or three centimetres back of the posterior boundary of the tumor. From

there the needle was carried at a right angle to the long axis of the tongue circularly around towards the ramus of the jaw, and then, penetrating through the mucous membrane of the floor of the mouth, was brought back into the oral cavity. In order to give the loop as immovable a position as possible around the muscular tongue, the right border of the tongue was included by the needle in bringing it out. The other half of the tongue was included in the same manner. The ligatures included in each case a little of the tissues on the other side of the median line; they were drawn together and knotted, the ends, which hung out of the mouth, serving to draw out the tongue. Thus the operation could be performed without loss of blood. x.

SYMPTOMS AND TREATMENT OF PSORIASIS UNIVERSALIS.—Kaposi (*Cbl. f. Chirurgie*, 1878, p. 406; from *Wien. Med. Wochens.*) depicts the appearances presented by psoriasis universalis where the efflorescence is interrupted by very small islands of sound skin or by none. Under these circumstances the affection may easily be mistaken, by those unaccustomed to the diagnosis of skin diseases, for pemphigus foliaceus, lichen ruber, or pityriasis rubra. Even specialists may sometimes confound this form of psoriasis with eczema universale. While until recently prognosis of such cases has always been unfavorable, Kaposi has of late succeeded, by the employment of cod-liver oil, of inunction, of baths, and of caoutchouc clothing, in obtaining very good results,—that is, of temporary cure. For instance, cases are on record where, after a course of cod-liver oil, terrible eruptions of eczema followed the psoriasis. In these cases care must be taken to stop the oil in due time. Continuous baths, also envelopment in bandages covered with ung. diachylon, prove useful in these cases. x.

INJECTION OF TINCTURE OF IODINE INTO THE KNEE-JOINT.—Orlow (*St. Petersb. Med. Wochens.*, 1878, No. 12) has in ten cases punctured the synovial sac, drawn off the fluid, and injected dilute tr. iodine (1 to 3 parts water). After a few moments a portion of the injected fluid was allowed to escape. The reaction was not severe, the patients were treated in the out-patient clinic, and needed no further treatment. Eight of the cases were of serous synovitis, and two of gonitis sero-purulenta. x.

UTERINE MYOMA—SUBACUTE PERITONITIS FOLLOWING SIMPLE VAGINAL EXAMINATION—DEATH.—A. Poncet (*Cbl. f. Chir.*, 1878, p. 391; from *Gaz. Méd. de Paris*) had under his care a woman suffering from a voluminous uterine myoma. A few hours after a simple vaginal examination had been made, the patient was attacked by peritonitis, of which she died within forty-eight hours. Notwithstanding the fact that the various symptoms of peritonitis, swollen abdomen, severe pain upon the slightest pressure, continual vomiting, etc., were present, the post-mortem examination showed no signs of inflammation, excepting some four to five hundred grammes of sero-sanguinolent fluid in the cavity. The author considers the case one of septic peritonitis. x.

PENETRATING WOUND OF ABDOMEN—RECOVERY.—Pierantoni gives the case of a girl of 18 who received a penetrating wound of the abdomen. The intestines protruded and were wounded, and the external wound bled considerably. The wound of the intestine was sewn up, and then the abdominal wound was closed. A few days later, meteorism, vomiting, etc. The wound came open and some masses of fæces escaped. After this the patient's progress was favorable, and she was quite well within twenty-six days from the operation. x.

PREVENTION OF HEMORRHAGE AFTER THE USE OF ESMARCH'S BANDAGE.—It is generally regarded as one of the chief drawbacks to the employment of Esmarch's bandage that hemorrhage, often of an exceedingly stubborn character, is so apt to occur after operation. M. Riedinger has sought to avoid this unfortunate complication by the use of the induced current. This appears to possess the property of arresting consecutive hemorrhage even when ischæmia has existed some time, a result which seems to depend upon the contraction of the muscles and of the vascular walls. R.'s procedure is as follows. Having finished the section of the tissues and tied all attainable vessels, he submits the wound to the influence of the induced current, the electrode being armed with a sponge, for some moments, and then, removing the tube of the Esmarch's apparatus, secures those vessels which continue open. His success in numerous cases has been very satisfactory.—*Le Mouvement Méd.*, 1878, p. 263; from *Arch. Méd. Belge*. x.

PHILADELPHIA
MEDICAL TIMES.

PHILADELPHIA, SEPTEMBER 14, 1878.

EDITORIAL.

MARINE HOSPITAL SERVICE.

THE report of the Supervising Surgeon-General of the Marine Hospital service, Dr. Woodworth, which is just at hand, contains much matter worthy of the study of every one interested in the hygiene of water transportation, or of sea or fresh-water ports. On the whole, it is an earnest, manly document, and its suggestions seem to be worthy of most careful consideration. We have one criticism, however, to make. In an official report, and especially in one discussing proposed measures of reform, the tone should be not so much that of advocacy as of calm, judicial weighing of advantages and disadvantages. Hence the utmost care should be taken that no incorrect assertions are made. The two most important measures proposed by Dr. Woodworth are—first, the erecting of marine hospitals in such cities as New York and Philadelphia, and the abolition of the time-honored practice of boarding sailors in civil hospitals; second, the introduction of physical examinations of merchant sailors. In the discussion of each of these measures there are statements concerning the very vital question,—that of expense,—which we believe can scarcely be sustained.

Thus, in regard to the first, we read (page 9), "This would not add to the cost of the service the amount of salaries of nineteen additional medical officers, as might at first be supposed, as, at ports where medical officers are not stationed, the pay for medical attendance is counted as part of the per diem rate, which embraces the cost of board, nursing, and medicines." We cannot speak for all the ports, but we know that in Philadelphia, and we presume

that in every large city of the nineteen ports, the seamen are treated in large civil hospitals, where there is no pay for medical attendance, and no interest upon the cost of erection; they are taken also at a rate which does not compensate the actual average outlay upon patients. The hospital has to be heated, lighted, and officered whether the sailor is or is not in its wards: hence it is considered economically wiser to receive the marine hospital protégé at a little above the actual cost of the food and medicine consumed than not to receive him at all,—a conclusion clinched usually by the earnest desire of the medical staff to have as wide a range of cases for study as possible.

Under these circumstances it is very obvious that erecting and sustaining hospitals in Philadelphia and similar places would add very materially to the expenses of the service, provided of course the food, etc., were as good in those hospitals as the sailors now receive. Again, in these same city hospitals are found large staffs of the most eminent men of the profession, devoted each to some special branch of medical science; a band whose collective wisdom and skill certainly must be much greater than that of the single marine hospital surgeon, who would unite in his own person all the medical and surgical knowledge and skill available in the proposed hospital.

It certainly is very clear that there are many disadvantages in the proposed change, and the report of Dr. Woodworth fails to make apparent any at all commensurate advantages other than the magnifying of the number and importance of the marine hospital service, and the supporting by government of an increased number of physicians,—the last, indeed, a result, in the present overcrowded state of the profession, devoutly to be prayed for.

The discussion of the second reform urged by Dr. Woodworth is chiefly from the pen of Dr. P. H. Bailhache. Some of the argu-

ments which he brings forward in favor of it are extremely forcible, but chiefest of them all is that which is derived from the greater safety to be secured for vessels with their living and lifeless freight. If it be true that "bad sailors more than bad ships are the causes of losses," then the instinct of self-preservation certainly demands that sailors should be carefully selected, whatever difficulties may lie in the way. These difficulties seem, however, underrated by Dr. Bailhache. To be trustworthy, this examination must be made at least every time a sailor ships, and so we understand the proposition to be. This being the case, does not the following quotation contain an error?

"No expense to the government, to the seaman, or to the vessel attaches to this measure. The machinery for its economic and successful management already exists. The Marine Hospital Service, through its medical officers, can readily assume and perform all the duties required in making such an examination," etc. Now, it is notorious that no marine hospital service at present exists at many of our large cities, and, if it be created, the physician who is occupied with the sick cannot spend his time examining recruits.

How many surgeons would have to be added to the service in New York City alone? It would certainly be unjust to make the hospital fund pay this expense,—especially as the legitimate hospital expenses already are in excess of the receipts, as is shown by the fact that the surplus with which the year 1875 was started has been reduced at the close of 1877 about fifty-five thousand dollars, or, in other words, that during the three years named the annual expenses of the service have exceeded the annual receipts, on an average, eleven thousand dollars.

There is one point of view which, we believe, is not touched in the report of the Supervising Surgeon,—the justice to the individual sailor. The soldier and the

government sailor undergo a rigorous examination when they enter the service, but they are not forced to give up their livelihood when their physical condition begins to be impaired. There is no adequate provision for the merchant sailor when disabled. Often he must pursue his avocation unto death, or become pauperized. Now the question arises exactly at what point of physical disablement has society, represented by the government, the right to step in and say, "Our safety requires that you should be debarred from following the sea. You are used up, and can go to the poor-house, or do the best you can for yourself." It is true that society practically carries out this policy in various dangerous trades, but it enforces judgment not openly, but through what seems to the victim inevitable physical necessity. Surely before the pitiless law crystallizes in black and white upon the statute-book, a much wider and deeper discussion should be had than is afforded by the report of the Supervising Surgeon.

CORRESPONDENCE.

LONDON LETTER.

THE chief matter of medical interest recently has been the annual meeting of the British Medical Association. This, the forty-sixth of the Association, and the third visit to Bath, was not expected to be a monster meeting; nor was it so. It was a moderate gathering numerically and intellectually, except Prof. MacKendrick's demonstrations of acoustics and Mr. Wheelhouse's address in surgery. Bath has seen its palmy days, and is beginning to present evidences of senile decay; yet its magnificent spring will always keep a certain number of people about it. Placed in an amphitheatre of hills, it is a warm, steaming place, and is a capital winter resort; but in August this does not constitute an attraction. Nevertheless, the meeting went steadily to work. The recent death of the outgoing president, Dr. Eaton Wilkinson, of Manchester, deprived us of the pleasure of seeing him vacate the chair in favor of his successor, Dr. Wilbraham Falconer, the well-known physician of Bath, and for years the respected treasurer of the Association. He contrasted the present condition of the Association with what it was when last there, in

1848. Then its transactions were over in two days, there were only 1400 members, and its income was only £1400. Now six days are taken up fully with its annual doings, it numbers 7500 associates, and its income last year was £11,500. (In 1848 it had only the guinea of each subscriber; now there are the advertisements in its journal as a source of revenue.) Out of its present receipts grants can be awarded for the purpose of encouraging scientific research; and Dr. Falconer referred to the work done by the late Prof. Hughes Bennett, by Professors Rutherford and MacKendrick, Dr. Braidwood and others. After reviewing the progress and present position of the Association, the president went on to describe Bath. He said it was a city which was resorted to for its social and educational advantages as well as for its notable spring. Recently remains of Roman architecture, in the form of a large bath two hundred and forty feet in length and one hundred and twenty in breadth, have been found, which attest the fact that at that time the spring at Bath was one of repute and distinction. Athelstan, Edgar, and Osric, Saxon kings, visited Bath and enriched it by gifts of gratitude for the benefits they had received therefrom. This spring is no mite of a thing: it discharges daily no less than 181,440 gallons; and the solid matter discharged in a year amounts to 420 tons, or about one ton and a quarter per diem. It comes up at about an average of 115° F., and is uncomfortably warm to the hand. There is a large and spacious pump-room, with an outpour as from a two-inch pipe, and this steaming fluid is drunk by the visitors in the morning fasting. This water is stimulant especially to the secretions, and acts on the bowels and kidneys. It is apt to produce headache if taken at all freely. In atonic gout the water is very efficacious, but is not suited for acute attacks. In chronic rheumatism it often achieves a cure where all other remedies have failed. There has been established a mineral water hospital for the treatment of gout, rheumatism, and paralysis. The fact, which was clinically observed, of paralysis being relieved by a course of Bath waters, is very interesting, showing as it does the association of paralysis with lithiasis and the condition of arteries and heart found therewith. At the Mineral Water Hospital the Sections on Medicine, Surgery, and Public Medicine were held, and in the room where the Medicine Section held its meeting there is an old picture of a physician and surgeon seeing patients with rheumatic gout, paralysis, and leprosy: the surgeon is feeling the paralytic's left hand very awkwardly, while the right arm hangs helplessly down. The peculiar pallid skin of chronic Bright's disease is very happily caught by the artist, and the man clearly has a clot in Broca's convolution. The thickened joints of a woman are also faithfully

depicted; and a little boy, with a shaven head and a fisherman's woollen cap on, has on his arms an eruption, but what form of lepra I do not know. I asked Mr. Jonathan Hutchinson about it, and he said he would look at it and tell me; but I have had no opportunity of seeing him since this conversation, so cannot in this letter say any more about it. Such, then, are the maladies for which Bath waters are in repute.

Then the secretary read the annual report; and a very satisfactory one it was. He announced that the invitation for the Association to visit Cork next year had been accepted, and Dr. O'Connor, of that city, had been elected president-elect. Seven hundred and odd new members had joined during the year. New branches had been formed; and the first colonial branch had been started in Jamaica. Grants had been voted for scientific research and for the systematic investigation, by a committee, of hydrophobia and rabies. Mr. Middlemore, of Birmingham, had generously given £500, the interest of which shall be awarded triennially for the best essay on the progress of ophthalmic medicine and surgery. A circular had been issued to every board of guardians, asking them to recognize the importance of registering all outbreaks of infectious disease and sending copies of reports to the Registration of Disease Committee. The Association had also taken the most active interest in the medical reform bill recently before Parliament. It is an unspeakable relief to know that this bill has been thrown out this year, and is not likely to raise its head again for some time; and in the interval possibly something like a decent comprehension of the matter may be developed; for, although medical reform has been talked about since the writer was a child, most men seem glad that the different prospected bills have perished *in ovo*.

On Wednesday morning the work went on briskly, the address in the Medicine Section being given by Dr. Goodridge, the senior physician to the hospital. He reviewed our present knowledge of those pyretic processes called "fevers," their causation, the changes set up in the organism, and the consequences thereof. If all this were worked out and known, probably the difficulties in the way of rational treatment would be much reduced. "The doctrine of despair" was to be reprobated, and passivity might well be exchanged for the most active investigation, with advantage. In the afternoon the Sections went to work, and in the Medicine Section a discussion as to the diagnosis and treatment of intestinal obstruction attracted a crowded audience. The only part of the discussion which had any claim to novelty was the advocacy of surgical interference not only in order to unloose twisted folds of gut or unravel a volvulus, but for the purpose of ascertaining exactly the nature and relations of other

obstructions, including malignant growths, and of giving relief by the performance of Amussat's or Littre's operation. The recent experience of ovariologists has shown that exploratory incisions could be made into the abdomen with comparative impunity, especially with antiseptic precautions. We are "not losing our respect for the peritoneum," as Mr. Wheelhouse said, next day, in his surgical address, but certainly we are losing our timidity about interfering with it. The introduction of this feature of a discussion on some special subject is of recent date, and though these discussions have so far brought out nothing original, they enable a large number of the profession to hear the latest words that have been spoken on the subject, by what falls from the various speakers. The ad interim report of the Hydrophobia Committee was then read by Dr. Gowers. In the Surgical Section the papers were all connected with the bones, hip-joint disease, and subcutaneous section or excision, and spinal curvature, especially as to the use of the plaster jacket therein. It must be gratifying to American surgery, and personally pleasing to Dr. L. Sayre, to know how general has been the adoption of his plan of treatment, and the success that has been achieved thereby. In the Obstetric Section a number of papers were read, but none of them were of such interest as to call for any allusion to them. In the Public Medicine Section the papers ran on the compulsory registration of infectious disease; but unanimity of opinion was not attained. The feature of the Physiological Section was the address on Physiological Acoustics, by Professor MacKendrick, of Glasgow, who has recently published an admirable text-book of physiology for the use of students. He pointed out how the increasing rapidity of air-vibrations produced a tone by the use of Helmholtz's siren, and showed how a noise differed from a musical tone, the latter being due to regular vibrations. He then demonstrated how the flame of a gas-jet could be shown by a revolving mirror to be a steady line of light till the note to which each flame corresponded was struck, and then the flame became a series of lambent tongues of light. The lucid exposition of the professor, and his admirable manual dexterity in making his experiments successful, evoked a general request that he would repeat his experiments next day, when a large audience warmly applauded his lecture.

In the evening there was the usual conversation, given by the mayor and citizens, and not by the president and the members of the Association, this year, which was largely attended. Of course there were the geological sections of the city and neighborhood, fossils, prints of Bath in its palmy days, the phonograph and telephone, and a large muster of the aborigines. The girls of the west of England are said to be "poor, proud,

and pretty." As to the two first, nothing may be affirmed; but certainly as to the third attribute there was no positive evidence; but that might be due to the large infusion of outsiders who have come for various reasons to reside at Bath. The original feature was a quantity of Bath waters in bottle,—the first step in an attempt to introduce them into general use. The attempt may be successful, but the water is heavily handicapped as a beverage. It is not palatable even when cold and aerated. It cannot compete with the continental waters either as a beverage on the one hand or as a purgative water on the other. It is not pleasant enough to drink, and most people will prefer something shorter as a purgative. It may be questioned whether patriotism, in its present enfeeblement, will procure a sale for Bath water, unless its reputed stimulant qualities can attain for it a demand among the temperance party.

The usual temperance breakfast, under the presidency of Mr. Samuel Bowley, came off next morning. About one hundred and fifty were present. It may be asserted confidently that such an attendance does not faithfully represent the attitude of the profession, and it is impossible to escape the conviction that the chief factor in such an attendance is that it saves a breakfast in the hotel bill. The next factor is the prospect of some amusement. This year there was not much of the latter, and the only thing said likely to live was the writer's remark, in recommending the temperance body not to oppose the physiologists by stating that alcohol was not burnt in the body, "that the food-value of alcohol taken as a beverage was scarcely worth contending for: that before a man could make a substantial meal of alcohol he would be blind drunk." These breakfasts are useful enough, but there is rather too much of the setting up of teetotalers, not always even medical men, to tell the profession what the right view of the alcohol question is, than a following out of the programme, viz., to ascertain the views of the profession on the subject of the use or abuse of alcohol. Nevertheless, the profession are always glad to see Mr. Bowley on these occasions.

On Thursday morning Mr. Wheelhouse gave the surgical address, which was in every way worthy of him and the Leeds school which he represents. He traced the gradual growth of modern surgery, using illustrative cases from his own experience or that of his colleagues, giving to them that generous credit which is not always accorded to colleagues who are also rivals in practice. It is simply impossible to follow this address, or attempt any abstract of it; but a brief allusion to the last case he gave may be permissible. A man, in crossing a fence with a scythe, got a wound of the buttock, the scar of which is still nine inches in length. The sciatic nerve was completely severed, the

limb was a flail, and he presented himself at the Leeds Infirmary to have the useless encumbrance removed. Dr. Clifford Allbutt found that the muscular susceptibility to the faradaic current was all but extinct. Mr. Wheelhouse cut down and found the nerve-ends two inches apart, with a large ganglion of nerve-matter, etc., on the proximal ending. This was removed, and an attempt was made to approximate the nerve-ends by longitudinal sutures. To achieve this the leg had to be flexed to the utmost, and the ankle strapped to the buttock. Then it became possible to get the ends together. First came back some sensibility, then some motor power, and now the man is able to work for a living, though the injured leg is not yet equal to its fellow.

There was little of interest in the Sections, the most entertaining matter being the discussion on syphilitic neuroses in the Medicine Section. In the evening the public dinner was held, at which were present Dr. Marion Sims, Dr. Daly, of Pittsburg, Dr. W. H. Pancoast, of Philadelphia, Dr. Cleaver, of Keokuk, Iowa, and Dr. J. J. McAngear, of Fort Madison, Iowa. Dr. Marion Sims and Dr. Daly replied for the toast of "The Guests."

On Friday morning the address in Forensic Medicine was given by Prof. Douglass MacLagan, of Edinburgh. After describing how much more forensic medicine was studied in Scotland than in England, he proceeded to contrast the criminal procedure of Scotland with that of England. He upheld the view that it was the duty of the state to prosecute, and he inveighed against the present system in England. In Scotland they had a public prosecutor, while in England the cause of death was ascertained, or attempted to be ascertained, by a coroner,—an officer, he said, of Plantagenet times, who was aided by a jury who were the neighbors of the injured person, and who therefore knew something about the cause of death. He did not approve the modern idea of medical coroners. He thought the lawyer on the bench and the doctor in the witness-box constituted the right men in the right place. He thought English jurisprudence might take a hint or two from the Scotch usefully. Meanwhile the Sections had been going on without much interest, except a paper by Dr. Bell, of Bradford, on wool-sorter's disease. This is a form of acute septicaemia, which may prove fatal in seventeen hours. It occurs when moist wool or alpaca is being sorted, and if the wet bales are allowed to dry, all danger is avoided. The septic particles seem to be inhaled, and, if death is not so rapid, lung-symptoms are developed along with the general condition of septicaemia.

In the afternoon the mayor gave a garden-party in his beautiful grounds on the top of one of the numerous high hills which surround Bath. Some exquisite iced coffee and cream was served out to the guests on their arrival, and after that there followed ices, champagne

cup, and other good things which suit the inner man at the later stages of a week like this. The evening closed by a concert at Colston Hall, Bristol, given by the Bristol doctors, where there was excellent music. A special train ran the medical freight back and forwards. This closed the meeting, and next day everybody was off, either on the excursions or otherwise. But a word or two may not be out of place about Bath in addition to what Dr. Falconer said. Bath is a staid town, and its chief characteristics are propriety and piety. A staidness is seen everywhere. There is religious gravity in the air, as there is moisture in the atmosphere. Its sombre terraces, its serious-looking houses, the numerous tablets in the Abbey, all tell of grave and earnest women: very genteel poverty flourishes there and rules the town council. Respectability and seriousness are the two leading indications of Bath, and this ought to be made the most of. Such concomitants of the spring ought not to be overlooked, then, and in their advertisements on the labels of the newly prepared bottles of Bath waters should be printed, in black type, "the religious watering-place." The people who would be at home at Bath would have little affinity for the light and thoughtless gossipers who congregate at Harrogate and Homburg, who are not only in the world but "of the world." Such being the case, it seemed specially unfortunate that this "city of the sea" should have been chosen as the place where the vote of the Association should be taken as to the admission of women members. If the advocates of their admission ventured to hope that the guardian spirits of Bath would be favorable to their cause and dishearten their opponents, they were very much mistaken. And though the speeches were not very convincing, and Mrs. Garrett Anderson made an eloquent and vigorous speech, the conclusion was evidently a foregone one: the opposition was overwhelming. The ladies have lost the privileges of the membership, and really one does not exactly know whether they should be congratulated or commiserated with: they at least have been saved, even if by their opponents, from being wearied by having to attend the Sections, or being bored by the reports of special committees, and still more from the terrible penance of listening to the after-dinner speeches of the annual banquet. Their lot may be a hard one, they may have been very badly treated; but there are those who think that they ought to draw a long breath of relief at the thought of what they have escaped.

The hospital at Bath, as also the Exeter Hospital, is a century old, of the days of palatial buildings, with the feelings of the architect the great matter for consideration, rather than the form of building best suited for checking the spread of pyæmia. Both hospitals are well ventilated, and kept very clean and free from hospital smell, and the patients looked

well fed and happy, as the denizens of these two rich sunny counties should be.

Instead of joining any of the excursions as usual, and leaving the British Medical Association to go its way, my steps were turned to Wonford House, Exeter, the asylum for the middle and upper classes of Devonshire, Cornwall, Dorset, and Somerset. It took its origin "at a meeting of the grand jury, held at the castle of Exeter, the 16th day of March, 1795, Sir Boucher Wrey, Bart., foreman, in the chair." A subscription was opened and a meeting held on the 29th of July, 1795. Some of the rules then passed contrast the treatment of the insane at that time with their present management very strongly. Rule 2 is, "No patient shall be chained or handcuffed without the previous knowledge and approbation of the physician or apothecary." It is possible to suspect, however, that general orders were often given by the physician or apothecary, for we read another rule, "The physician and apothecary shall visit the asylum every Tuesday morning at twelve of clock, and oftener when any urgent case shall require peculiar care and attention." Now there is a trained medical superintendent resident, and there is not a strait-waistcoat, even, on the premises. Other measures than brutal violence are now found much more efficacious. There is another rule which has an ominous look about it; it runs thus: "that the feet of every patient under confinement or in straw be carefully examined and rubbed and covered with flannel every night and morning during the winter months." This calls up some painful suspicions as to what refractory and excitable patients had to go through in the days of old. Now such things are unknown, even in pauper asylums, to say nothing of the denizens of Wonford House. A beautiful three-story building of ornamental character, it stands about a mile from Exeter, with a range of hills across the Exe facing its front. In its grounds are targets for archery, and hoops for croquet, and the attendants and patients at cricket enjoy themselves famously in the balmy air of South Devon. Inside are drawing-rooms, music, billiards, every home comfort that can make the patient feel the restraint as little as possible. But still beneath this silken glove there is the iron hand. Sitting with a grand old parson, mad as a hatter, but a gentleman withal, I tried to throw the window a little farther up: it successfully resisted my efforts, when the old boy, with a chuckle of much inward satisfaction, said, "Don't you remember you are in a mad-house?" Those windows have nothing repulsive about them, but the suicidal patient has no chance of getting out either of the house or his troubles by the open window. Then the superintendent is a cultured gentleman, who sympathizes with every one of these mad folks and makes their captivity as little galling to them as is possible. We went out

for a drive the other day, with two or three of the poorer patients, whose friends cannot afford drives, with us. There may be a strain of insanity in me, but the insane are far more interesting and entertaining to me than the sane,—infinitely more so. And if ever this latent strain develops into something more pronounced, I trust some connection with the four western counties will be established, in order that Wonford House may be my home during the period of my mental alienation. With these gentle, pleasant Devon patients to associate with, and this thoughtful, able doctor to look after me, insanity and captivity are robbed of half their terrors. In such a place the insane are infinitely better off than at home, even if surrounded by luxuries and kind and sympathizing friends. Gentlefolks carry their training and their evidences of good breeding with them even beyond the portals of an asylum; and the courteous Devonshire people are amiable and lovable even in their madness. Their character harmonizes with their pleasant landscapes, their balmy temperature, their sunny skies, for Devonshire is indeed "the garden of England;" while the beauty of the girls, with their dark locks, their bright eyes, and clear complexions, is linked with a stalwart bearing, manly yet gentle, of the men,—the men truly of Charles Kingsley's "Westward Ho," the crew of the *Rose*. As I sit at the open window, with the breeze rustling amidst paper and fluttering leaves both of open books and ferns, the prominent feeling is that this place is as near Paradise as man may hope for in his unglorified state. A well-conducted lunatic is the most agreeable of all companions, when the mental aberration is just sufficiently pronounced to give piquancy to the character. The sylvan beauty of the scene, the clear blue sky flecked with white clouds, the inspiring breeze, all conduce to the feeling of inward satisfaction, and raise high the anticipations of pleasure to be derived from an excursion up the Dart with some of these interesting patients as companions. But my letter must come to an end, as these personal details can scarcely be interesting to readers in search of medical details. One item may be pardoned. This morning, as the doctor read the record of his tell-tale clock, his face was clouded for a moment. On inquiring if anything was amiss, he smiled, and said, "No: this is the clock's fault." It must be a very clever tell-tale clock that can indicate its own shortcomings!

J. MILNER FOTHERGILL.

TO THE EDITOR OF THE PHILA. MEDICAL TIMES:

DEAR SIR,—At a recent meeting of the New York Academy of Medicine, Dr. Lewis S. Pilcher read a paper upon the subject of Colles' fracture, the chief point of interest being the theory which he advanced as to the mechanism of the production of this injury.

He rejected the commonly received explanation that the radius, being caught between two counter-forces, gives way at the weakest point, about one-half inch from its lower extremity, but asserted that the force immediately concerned in producing the fracture acted through the medium of the anterior radio-carpal ligament. Forcible extension of the hand being made, this ligament becomes tense, and if the strain is continued, either the ligament must give way or the radius, which is in the position of a bent lever of the first class,—the fulcrum being the posterior surfaces of the scaphoid and semilunar bones,—must suffer fracture.

This was followed by an exhibition of some specimens in which the fracture had been produced in the above manner.

In the discussion which ensued, Dr. Frank H. Hamilton remarked that he was "proud to know that it was an American surgeon who had made the observations, and felt highly honored that the New York Academy of Medicine was among the first medical societies to which these observations had been unfolded." Dr. A. C. Post suggested "that the fracture should be called Pilcher's fracture instead of Colles' fracture." Dr. Willard Parker had "listened with profound interest, and had been much instructed by Dr. Pilcher's paper," etc.

As these distinguished surgeons estimate so highly the value of this explanation of this common accident, it may be interesting to note that substantially the same theory has been taught for years at the University of Pennsylvania by a Philadelphia surgeon. In my notes on the lectures of Prof. D. Hayes Agnew I find that he defines Colles' fracture to be "a transverse fracture of the radius occurring just above the articulation," and says that while "Barton's fracture—or a fracture extending into the joint—is caused by direct pressure of the carpal bones on the posterior rim of the articulating surface of the radius," yet that the conditions in Colles' fracture are different. "It falls on the hand when it is strongly extended, the posterior—not the articular—surfaces of the scaphoid and semilunar bones are driven up against the radius and form a fulcrum. The anterior ligament (radio-carpal) is brought up strongly and with great tension against the front of the articulation, and as it (the ligament) usually resists the force, the solution of continuity takes place at the point to which the strain is transmitted, which is the lower end of the radius, a short distance above the joint."

This explanation is also that given by Dr. Agnew in his "Treatise on Surgery," the first volume of which is now in press and will be shortly forthcoming. If there is any credit in priority in these observations, and if there is any reason for a particular community to feel honored by them, the credit and the honor belong here and not in New York.

Yours truly, J. WM. WHITE.

222 SOUTH SIXTEENTH STREET, August, 1878.

PROCEEDINGS OF SOCIETIES.

PATHOLOGICAL SOCIETY OF PHILADELPHIA.

THURSDAY EVENING, JUNE 13, 1878.

THE PRESIDENT, DR. H. LENOX HODGE, in the chair.

Cancer of the right kidney—Thrombosis of ascending cava and left renal vein. Presented by Dr. JOHN GUITÉRAS.

HENRY F., a German, æt. 68. Admitted to the Philadelphia Hospital the 27th February, 1878. His legs were swollen, hard, shining. Eight or ten ulcers, about the size of dimes, could be counted on each limb. They were surrounded by a dark purplish skin, showing no tendency to heal. The swelling extended up beyond the popliteal spaces. It was more marked in the left leg. He was anæmic, short of breath, without any thoracic lesion apparent. The gums were spongy, the teeth loose, his diet bad, his residence the out wards of the hospital, where scurvy abounds. As a case of scurvy the man was treated, and improved much. He gained in vigor, in flesh, and in color. The ulcers healed, leaving some indurations. The right leg returned to a normal size, and the left was almost as much reduced. Now it was found that the horizontal position was necessary to keep the legs in this condition. If he stayed up they were sure to swell. Outside of this, and an occasional accident that will be mentioned shortly, he appeared in very good health, had very good color, and was the most cheerful patient in the ward; had no pain whatever; no albumen; no casts in the urine. The belly was large, but nothing that might not be normal. The liver was large, and the superficial abdominal veins were somewhat distended. The heart was fatty, though not to a marked degree. An anæmic murmur at the base disappeared under treatment. While in the wards he had two severe attacks of hæmaturia, relieved by ergot and gallic acid. They left no trace behind.

The third and last attack commenced on the 20th of May. He was in bed, getting over a slight swelling of the legs, brought on by sitting up for two days. I did not see him at this time, but I gather from the notes of Dr. Roberts, the resident physician, that on that day the urine became dark and clotted in the vessel. On the 21st, "Passed no urine in twenty-four hours; bladder rose high up in the hypogastrium; passed a No. 20 catheter, and removed, by aspiration, about two ounces of clots. This was followed by the discharge of about fourteen ounces of bloody urine. In spite of astringents and hypodermics of ergotin the hæmaturia continued. On the 23d the hemorrhage was so

great, and the condition of the patient such that, as a last recourse, Dr. Bruen suggested the injection of a weak solution of Monsel. The hemorrhage was controlled, but the weak pulse did not improve, and he soon commenced to vomit. He continued to pass very small amounts of dark-colored urine."

May 26.—I saw him, and found distinct symptoms of uræmia. He lay unconscious, with stertorous breathing, and occasional twitching of the right side, face, and arm. With the right hand he continually scratched the chest and abdomen. The left side seemed to be paralyzed. There was rotation of both eyes, and slight turning of the head to the right. The face was not engorged. The pulse had become slower and full. The legs were not more swollen, but the superficial abdominal veins were more distended than usually. The temperature was subnormal. He died the same day.

The post-mortem examination revealed a general infiltration of the right kidney, with encephaloid cancer. The organ was uniformly enlarged to the size of a foetal head. There was no cancerous growth elsewhere. The vesical plexus of veins contained coagulated blood. A yellowish, firm, partially-adherent coagulum extended up the internal iliacs and the cava to the level of the renal veins. A firm, loose plug was found obstructing the left renal vein. The brain was anæmic and œdematous. There was no marked atheroma, no hemorrhage, no embolus, no thrombosis.

Remarks.—I am sorry that a more careful examination was never made of this patient. His general condition excluded all idea of abdominal cancer, though the persistent swelling of the feet pointed in this direction, together with the hæmaturia. The theory of scurvy seemed to explain all the symptoms, and an examination of the abdomen was neglected, which would certainly have led to the discovery of the tumor.

I thought that an occlusion of the cava and renal veins was the only condition that could give rise to the symptoms present before death, especially after taking into consideration the injection of Monsel into the bladder. Of course I expected to find that the function of both kidneys had been interfered with by the clot. I also expected to find some obstruction in the cerebral vessels.

In the discussion on the diagnosis of renal enlargement, Dr. Guitéras said that his own and the other cases presented should be used as evidences, not of the impossibility of detecting renal enlargement, but of our neglect to look for it. He was sure that in the great majority of these cases the region of the kidneys had not been examined. He remembered a case where Dr. Pepper had diagnosed, by physical exploration, a nephritic abscess in a case where the symptoms could be well accounted for by an empyema that was also

present on the same side. He thought that palpation gave valuable information as to the size of the kidney. He had found that by frequent percussion he could always satisfactorily map out the organ. Very often in the first examination no results could be obtained, but subsequently one became able to recognize the changes in the conditions of collapse and inflation of the surrounding guts that act as obstacles to the examination.

Colloid cancer of the abdomen. Presented by Dr. THOS. M. DRYSDALE.

I visited Mrs. Margaret M. Rice, in consultation with Dr. William B. Atkinson, October 30, 1877, to examine a tumor of the abdomen, which was supposed to be ovarian. She gave me the following history of her case. She was thirty-three years old; had always been healthy, except an attack which her medical attendant called "bilious intermittent fever, with liver complaint." This was twelve and a half years since.

She first menstruated when between sixteen and seventeen years old, and had no pain nor trouble of any kind; has always been regular, except when pregnant, until two months ago, when she missed it, and has not menstruated since. Her menses were never accompanied with pain, were normal in amount, and usually lasted two and a half or three days.

She was married when twenty-one years of age, has had three children, all now living, the youngest three years old last May, the next six next month, and the oldest nine the 1st of October. Her labors were perfectly natural and easy. The first was slow, the others quick. She nursed all her children. After marriage she was always troubled with leucorrhœa, often quite profuse.

She has been losing flesh for over a year, and has become greatly emaciated and anæmic. She has had no trouble with her bladder; her bowels are natural in every way; her appetite remains good, and up to this time she has done her own work.

The enlargement was first noticed two years ago, in November, when nursing her child, and just at that time she had an attack of nervous prostration, which her doctor told her was owing to prolonged lactation, and advised her to wean her babe.

The swelling commenced in the centre of the lower part of the abdomen. It was comparatively small till last June, and then rapidly increased, causing her to look "just like a woman in the family way." This rapid growth has continued to the present time.

She has never had pain in the abdomen until two months ago, and then for a short time had severe shooting pains, which were supposed to be owing to an umbilical hernia. She was first attended by Dr. Ruttey, who used electricity. She then went to the University Hospital, where she was examined by

Dr. Goodell and told she had an ovarian tumor. Dr. William B. Atkinson was next called upon, and he, in company with Dr. I. Williamson, tapped her, and obtained about a quart of a jelly-like matter. These gentlemen also diagnosed an ovarian tumor.

I found the abdomen very prominent and enormously distended. A large umbilical hernia existed, filled with fluid, a contraction in the middle causing it to appear double. The abdomen was dull on percussion, except along the region of the ascending colon. The right flank was resonant, the left dull, and along the edge of the ilium semi-resonant. Fluctuation could be detected everywhere, except in the right hypochondriac region. The tumor was smooth, elastic, and immovable. There was neither pain nor tenderness on pressure.

Examined per vaginam, the uterus could be felt low down, with the cervix pushed towards the right side, and pointing that way. The uterus was immovable. The sound entered three inches, and so anteriorly that the handle had to be pressed against the perineum to insert it. The back of the pelvis was filled with hard, fixed masses, which felt like uterine fibroids, and were tender to the touch. The uterus was intimately connected with or imbedded in these growths.

December 8, 1877.—In company with Dr. W. L. Atlee and Dr. Atkinson, I tapped her, and removed about three quarts of a colloid matter, which resembled in color, consistence, and general appearance calf's-foot jelly. There also escaped small quantities of a thin greenish fluid. It required fully an hour to obtain this quantity, although a large canula and trocar were used.

After tapping, the epigastric region and whole upper part of the abdomen became resonant on percussion. On deep pressure, a hard mass could be felt in the left hypochondrium. Palpation gave the sensation as if a stratum of thin fluid existed between the abdominal wall and tumor. The mass could now be moved to a certain extent. My diagnosis was given as colloid cancer of the abdomen, and an operation declined.

The tapping relieved her of the feeling of distention, and on visiting her the next day I found her sitting up. Her color had improved, her lips were red, and she said she felt quite comfortable. Dr. Atkinson now took entire charge of her case. She continued to emaciate, the abdomen enlarged, and in a month or two the umbilicus ulcerated, an opening forming, through which large quantities of colloid substance, and a greenish fluid, were discharged, as much as a bucketful of the last escaping at one time. When I last saw her, June 5, the abdomen had entirely changed in appearance; the lower part had retreated, more especially on the right side, leaving a hard mass filling the entire upper part of the belly, and another

and softer mass the left lower portion. She died June 8, 1878.

Assisted by Dr. W. L. Atlee, Jr., I made an examination of the body June 9. The emaciation was extreme. The abdominal wall was so thin that my first incision penetrated the cavity. The peritoneum was thickened and opaque; its surface was unevenly studded with white deposits, resembling the diphtheritic membrane; the subperitoneal areolar tissue was converted into colloid, forming a layer from an eighth to a quarter of an inch in thickness. There was no organ which had a natural appearance. Filling the epigastric, the right and left hypochondriac, and left lumbar regions was an enormous colloid tumor, which careful examination proved to be the degenerated omentum. When cut, it had the appearance of a honeycomb, the cells or alveoli being filled with a pale-yellow jelly. Below this mass were the small intestines, lying empty, and close to the back. Below these again was a large soft mass filling the hypogastric and right inguinal regions, and stretching far towards the left. When the adhesions of this were detached, it was found to be the right ovary in a state of colloid degeneration. The uterus, which was found below this mass, appeared healthy on superficial examination. The left ovary had degenerated into a single cyst, filled with jelly, which not only occupied the upper portion of the pelvis, but sprang up as a globular tumor behind the uterus. The contents of the abdomen were incorporated into one tumor by this colloid material, and scattered through the gelatinous body could be found masses, varying in size, of the more fibrous or honeycomb-like material. It was a collection of these that were felt during life surrounding the uterus.

I present the specimens, but will not describe them minutely, both to save time, and because the general and microscopic appearance of this form of cancer has been so ably treated of by Sir James Paget in his *Lectures on Surgical Pathology*, to which I would refer you (p. 772). I would merely say that all the specimens I have examined of this disease correspond closely with his description, except, in two instances, the absence of the nucleated cells, which he (after Lebert) describes as lying free in the colloid substance.

The object which I had more particularly in view in bringing these specimens to your notice was to point out the great similarity of symptoms between this form of disease and that of multilocular ovarian tumor. In fact, so closely do they resemble each other that I believe it impossible to distinguish between them during life, except by tapping. This case was diagnosed as ovarian tumor by more than one able physician, and I certainly should have agreed with them before the tapping was performed, but, having met with

these cases before in the practice of Dr. Washington L. Atlee, I was familiar with the jelly-like matter, and so saved from an erroneous diagnosis.

Dr. Atlee, in his great experience of abdominal tumors, has met with but six of these cases, and they were so much alike that a description of one would apply to all, and in nearly all were they mistaken for ovarian disease by their physicians until the tapping revealed their true nature. This colloid or jelly-like material is unmistakable, and once seen can be always recognized.

Case of recurrent fibroid of axilla. Presented by Dr. NANCREDE for Dr. J. ASHHURST, JR.

The tumor was removed from Claude Conrad, aged 21 years, who was admitted, October 11, 1876, to the University College Hospital for encephaloid of the axillary, which diagnosis was revised to sarcoma. Professor Agnew removed the growth on the above-mentioned date, and he was discharged, cured, October 31, 1878. Since that time five other small recurrent tumors have been removed by his family physician in Factoryville. On May 28, 1878, Dr. J. Ashhurst, Jr., removed the present tumor, with two smaller growths, making in all eight tumors removed since the first was excised by Dr. Agnew. Macroscopically, this growth resembles strongly a uterine fibroid, but microscopically was seen to consist of numerous cells of a connective-tissue type, imbedded in a hyaline intercellular substance, presenting no appearance of fibrillation. Some of the cells were becoming decidedly spindle-shaped, evidently developing into fibre-cells.

THURSDAY EVENING, JUNE 27, 1878.

THE PRESIDENT, DR. H. LENOX HODGE, in the chair.

Three cases of heart disease associated with kidney disease. Presented by Dr. JAMES TYSON.

GEORGE C., colored, age 45, native of Philadelphia, occupation barber, was admitted to the Philadelphia Hospital, April 5, 1878, in a very feeble and prostrated condition, frequent pulse, general œdema, and great dyspnœa. He was a very large, stout man. Although his intelligence was perfect, he never was inclined to answer questions. He said he had been healthy previous to his present illness, which had been of but a few weeks' duration; he denied having had rheumatism or syphilis, or having used liquors in excess. Bowels constipated; tongue coated; no appetite; great thirst. Cough with white frothy expectoration. Feet very much swollen; whole body œdematous. Skin dry. Pulse 120, dicrotic. Urine diminished in quantity, and highly colored.

Physical examination.—Area of heart's dul-

ness much increased; systolic murmur at the base; later, also, a mitral systolic murmur was observed. There were moist râles all over the chest and back. Examination of urine revealed albumen in variable quantity at different times, from one-quarter of bulk examined, to a trace; no tube-casts, but often much epithelium from the kidneys, and a great excess of urates.

Diagnosis.—Aortic stenosis and slight mitral insufficiency. He improved for a short time under digitalis and tonics. A laborandi sweat did not act favorably. Soon his condition grew worse; œdema became more marked; and the patient was compelled to remain in sitting position during the last two weeks on account of excessive dyspnœa. He died May 12, 1878.

Autopsy.—Heart hypertrophied and dilated. Aortic valves slightly atheromatous and stiffened, also fenestrated, but not thickened. Mitral cusps evidently, but not markedly, thickened. Aorta dilated, and highly atheromatous.

Lungs œdematous; some pleuritic adhesion on the left side.

Kidneys enlarged and swollen; in a state of parenchymatous nephritis.

Liver and spleen normal.

Brain not examined.

Remarks.—The most interesting facts in the history of this case are the slight degree of valvular lesion compared with the extreme degree of the symptoms, the marked dyspnœa and excessive dropsy, and the fact that there was also chronic tubal disease of the kidneys, with very few casts.

J. M., colored, age 56, native of Virginia, was admitted to the Philadelphia Hospital four times. First admission in May, 1877, for one week, with the diagnosis of aortic regurgitation. Second admission in July, 1877, when he was transferred to the venereal wards. Third admission in December, 1877, when he remained in the medical wards until March last, under the care of Dr. Guitéras; diagnosis, chronic Bright's disease. Last admission, May 24, 1878, under my own care. When I saw him he was in a very feeble, prostrated condition, frequent pulse, great dyspnœa, and not able to speak.

Physical examination.—Dulness on percussion over the right side of chest; region of dulness changed by change of position of patient. The heart was pushed much to the left side. First sound of heart reduplicated; slight systolic murmur at the base, and also a mitral regurgitant murmur. Dr. Guitéras also found marked dulness over the upper part of sternum; right pulse larger than left.

Examination of urine.—Albumen only a trace; urates in excess; hyaline and some granular tube-casts.

Diagnosis.—Pleuritic effusion; hypertrophy of heart; aortic stenosis and mitral insufficiency; interstitial nephritis. Dr. Guitéras

diagnosed, in addition to this, athroma and dilatation of the aorta.

Died on June 9.

The *post mortem* notes read as follows:

Heart.—Hypertrophy without dilatation; walls of left ventricle one inch thick; posterior leaflets of mitral valves thickened, the chordæ tendinæ of left valve much thickened, pulling out the muscoli papillaris, which were extremely hypertrophied and unusually hard. Semilunar aortic valves slightly thickened, and atheromatous. Aorta greatly thickened, and generally atheromatous, and *dilated*; about one inch from its commencement it became double.

Lungs.—Right lung much engorged and congested, the lower lobe completely consolidated, and entire hemorrhagic infarction. Extreme pleuritic effusion on right side.

Kidneys contracted, not more than three inches long; granular.

Liver, intestines, and peritoneum healthy.

Brain not examined.

Microscopic examination of kidneys confirms an advanced interstitial nephritis.

George J., white, age 35, native of Philadelphia, a laborer, admitted to the Philadelphia Hospital, April 17, 1878. He was a large, strong man, has never been sick, except an attack of rheumatism, which he had in 1871, for a few weeks, and never since. He had syphilis, and has been also a drinking man. His sickness dates only from February last, when he took cold, being one day whilst perspiring very much exposed to snow and cold wind. Had cough, fever, shortness of breath, and pain in the region of the heart. A few days before his admission to the hospital his feet began to swell, the swelling extending later over the whole lower extremities and the scrotum. He did not complain of any pain, but had very great dyspnœa. Appetite good; excessive thirst; quantity of urine diminished—only a trace of albumen; no tube-casts; some epithelial cells from the kidneys, and some blood-corpuscles.

Physical examination.—Area of heart's dullness increased. Marked systolic murmur at the base of heart, and a murmur is heard in the right carotid; a mitral murmur was also observed coincident with the impulse, and transmitted to the axilla. Vocal fremitus diminished at the left side of chest.

Diagnosis.—Mitral insufficiency and aortic stenosis, and chronic pleurisy.

The patient died suddenly and unexpectedly, May 12.

Autopsy.—*Heart* much hypertrophied; cavities much thickened; chicken-fat clot in right auricle. Extensive vegetations on aortic sigmoids, with partial calcareous degeneration of the same. Marked thickening of mitral valve.

Lungs.—Hypostatic congestion of both lungs; pleuritic adhesions on left side.

Kidneys large, swollen; capsule strips off

easily. Left kidney with marked hemorrhagic infarctions; in both kidneys minute hemorrhages over the surface. Cortex increased. Microscopic examination reveals a parenchymatous nephritis, with capillary congestion and extravasation of blood.

Dr. GUITÉRAS said that at a previous meeting he had referred to several cases where he had succeeded in diagnosing slight dilatation of the aorta by means of respiratory percussion. In some of the cases the diagnoses have been since that time confirmed by autopsies. This had led him to attach much importance to the systematic use of this method of percussion for examination of the aortic region. He made it a rule never to examine this region without percussing at the end of a forced expiration. As the lungs collapse over the large blood-vessels, it is surprising to find how clearly the area of dullness of a dilated aorta becomes apparent.

The first case he had examined after this manner was one where aneurism had never been suspected, and where no dullness could be obtained by ordinary percussion, yet he was enabled to demonstrate to a large class an area of almost flatness over the right border of the sternum, above the third rib. Slight pulsation could also be seen towards the end of expiration. Other symptoms have developed since, leaving no doubt as to the diagnosis.

In Case No. II., a colored woman, 38 years old, the symptoms of pressure on the trachea were well marked. Symptoms of vascular disease rendered it probable that the source of pressure was an aneurism, but the dullness at the upper piece of the sternum could only be detected at the end of forced expiration.

In Case No. III., an old woman, there were evidences of valvular disease, both mitral and aortic. The aortic systolic murmur was so feeble (weak ventricle) that the existence of lesion of the semilunar valves had been doubted. By respiratory percussion the presence of aortic dilatation was demonstrated, and the diagnosis was subsequently confirmed at the post-mortem.

Of Case No. IV. no notes had been kept. During life the enlargement of the vessel could only be demonstrated at the end of expiration. Diagnosis confirmed by the autopsy.

Case No. V. was the one now before the Society. When this case was admitted early this year in the wards of Dr. G. at the Philadelphia Hospital, he supposed it to be one of contracted kidney with cardiac hypertrophy. But his attention was called to examine the aortic region by a prominence of the right sterno-clavicular articulation. This had nothing to do with the aortic lesion, but pulsation at the base of the neck was noticed, with inequality of the radial pulses. Respiratory percussion led to the discovery of a well-marked area of dullness.

Case No. VI. was examined previous to these investigations, but it is mentioned because though the dilatation was not greater than in some of the cases just described, yet it could be detected by ordinary percussion. The emaciation of the patient probably accounts for this. She died of cancer of the uterus.

Case No. VII. was seen only yesterday, after a prolonged absence in Cuba. In 1876 Dr. G. had found evidences of disease of the aortic valves and of atheroma of the arteries in its early stage. During the same year he saw him, in consultation with Drs. Alonzo Clark and Austin Flint, of New York, and Dr. Landeta, of Havana. Dr. Flint confirmed the diagnosis, but Drs. Clark and Landeta suspected the existence of some dilatation of the arch. Dr. Clark based his diagnosis upon evidences obtained by ordinary percussion. Now the patient presents all the symptoms of aneurism of the ascending portion of the arch. It is very probable that respiratory percussion would have made the lesion more apparent in its incipency. Curiously enough, at present, when there can scarcely be any doubt of the existence of an aneurism, the percussion resonance over the aortic region is but very slightly impaired, and is not influenced by respiration. This is due, Dr. G. believes, to adhesions fixing the lung.

He also desired to call attention to the value of pulsation of the neck, as being a very conspicuous symptom in cases of aortic disease, both of the valves and vessel. It should always lead to a careful examination of the aortic region. In differences between the carotid pulses and in unilateral thrill pretty strong evidence is found of disease of the aortic arch. Of seven cases presented, five had unequal radial pulses. In No. II. this symptom was absent; in No. IV. it was not looked for.*

Unfortunately, the specimen of double aorta was removed from the body before the anomaly was noticed. At the first glance it was difficult to determine whether the double aorta was not a dissecting aneurism. There was a slit about an inch beyond the left subclavian artery leading into the accessory aorta. The latter ran up to the subclavian, forming a pouch, and downwards as far as the mutilated specimen shows. Further examination leaves no doubt as to this point: the edges of the slit are smooth, the lining membrane of the anomalous channel has the normal appearance, and, most important of all, the orifices of the left intercostal arteries are seen in the outer wall of the false aorta; the right intercostals arise from the main vessel near the septum; in the latter some aborted orifices may be found.

* All these cases will be published more in detail, together with observations on the normal respiratory percussion of the aortic region.

As a point bearing on the pathogenesis of atheroma, it may be mentioned that, whilst the arch of the aorta is well dilated, and, together with the rest of the main channel, is highly atheromatous, yet in the accessory aorta, where the blood-pressure must have been much less, there is scarcely any atheroma.

Cancer of the ascending colon. Presented by
R. G. CURTIN.

John McL., æt. 64 years, married, native of Scotland.

The patient, who was by occupation a stone-cutter, came under observation on the 15th of January, 1877.

When he was 21 years old, he had an attack similar to the one for which he came under treatment last January, except that he did not then notice any tumor at the seat of pain. This seizure passed by, but kept him so reduced that he was for a time unable to work.

In the early part of 1876 he had some rheumatic symptoms in the legs, and suffered from weakness and vertigo.

With these exceptions the patient stated that he had been perfectly well up to within eight months of the time he applied for treatment.

In the month of June, 1876, while lifting a heavy stone, he suddenly felt a stinging or *jagging* sensation, of a warm or burning character, in the right iliac fossa, which continued for two weeks, sufficiently severe to unfit him for work.

About two months after this lift, his attention was directed for the first time to a tumefaction in the right iliac fossa, at the point in which he had felt the sensation of pain. This tumor was somewhat tender on pressure, becoming more so as it increased in size. The patient compared the pain to that of an abscess, as if matter were present.

In August he had a great tendency to nausea, and frequently vomited. For a month or two only he belched up "wind and slime." When swallowing food he noticed an inclination to eructation. He would feel the gas working up and down, and obtained temporary relief after belching.

Never passed blood with his stools. Bowels were regularly opened until about October, 1876, when first symptoms of constipation appeared.

He would go to stool with flatulent colicky pains, and fail to be relieved. Suffered much from *rumbling* in the bowels, accompanied with pain.

When first seen, in January, 1877, he had coughed for two months, expectorating about a tablespoonful in twenty-four hours.

Auscultation revealed, on the left side, noisy and harsh inspiration, with audible and prolonged expiration, about the mammary region; on the right side in the correspond-

ing region harsh and noisy inspiration, with expiration same as on the left side.

There was progressive loss of flesh and strength, and the patient died, on the 18th of May (last), of exhaustion and starvation. There was extreme emaciation.

Post-mortem.—On opening the abdomen, a tumor, the size of the fist, was found about one-third of the way up the ascending colon. The only evidence existing of peritonitis was a small patch outside the tumor, where a loop of the upper part of the ascending colon was bent down and attached to the tumor. The tumor was hard, and on section the tissue grated under the knife like cartilage. The inside of the tumor was in an ulcerated state, with patches of red, pulpy mucous membrane scattered here and there. The alimentary canal was patulous through the tumor, but very much constricted.

The difficulty in making a diagnosis in this case from chronic perityphlitis is apparent from several considerations. First, he had stated that he had had a soreness, with evidence of inflammation, in the same place when a boy. Second, the soreness came on suddenly after a heavy lift. The growth was very slow, and ceased a year before death.

A daughter of this man had an ulcerated condition of the small intestine, seated near the same place, two years ago. The specimen was presented to this Society.

Dr. Seiler made a microscopic examination of the tumor, and found the muscular tissue of the intestine at the place of the new formation to have disappeared; in its place a connective tissue forming alveoli, in which were epithelial cells, was found.

Dr. Tyson said in reference to one of the symptoms mentioned by Dr. Curtin, the sensation "as if something had suddenly given way" within the abdominal cavity, that he thought this symptom was due here to the rupture of an adhesion, which had formed at a previous period, when inflammation first set in, due to the irritation of the growing mass. Some years ago he had met the same symptom in the case of a young man, who, while gunning, felt something give way in the abdomen as he made a long leap; immediately peritonitis was developed, and he died promptly.

Embolic pneumonia. Verbal communication by Dr. GUITÉRAS.

Dr. GUITÉRAS desired to relate very briefly a case bearing upon the discussion of embolism. He had found at the Philadelphia Hospital a pair of lungs presenting numerous small patches of gangrene. He had immediately suspected the existence of some focus of infection, since the tissue between the foci was healthy. On asking, it was discovered that the man had had during his rather short illness a dark and fetid discharge from one ear. The body was placed again upon the

table, and the skull was opened. A portion of the petrous portion of the temporal bone was found broken down, and the lateral sinus near it in a condition of gangrene; its walls and the contained clot presented a greenish hue, and were fetid. The body had not commenced to decompose. The emboli washed from this focus were evidently infectious, and they gave rise to a process similar to the one taking place at the fountain-head. The patient was a robust adult.

REVIEWS AND BOOK NOTICES.

HANDBOOK OF OPHTHALMOLOGY. By Prof. C. SCHWEIGGER, of the University of Berlin. Translated from the Third German Edition by PORTER FARLEY, M.D., of Rochester, New York. J. B. Lippincott & Co.

It has not been so very long since medical science passed through the dangerous crisis in which a vague and fantastic speculation threatened to carry all before it. The general necessity of placing upon a firm basis the educational standard of a proper scientific method of obtaining knowledge, only by close and careful observation and study, has driven away empiricism. This has become evident in the several branches of the animal functions and economy, by the new ideas already opened, and the lucidity given to many intricate points hitherto considered obscure.

The science of medicine is one of practical truths; therefore its progress, to be near the truth, must be kept far removed from all speculation which is not well supported by immediate observation and experience.

Until within a short time the attention of students has been drawn, in relation to medicine and surgery, to general matters only; while the special parts have been neglected, or rather skipped over. But, thanks to a divine order, the development has been so great that the different parts and organs of the human body have undergone especial observation, with the result of calling attention to the necessity of their separation into different branches for particular study, thus giving rise to specialties in the practice of our profession. In none of the branches has there been more activity in the development of really scientific and practical results than in that of ophthalmology. For the past twenty-five years it has been the great study throughout Europe and the United States, and new brochures and books upon subjects relating thereto are continually issuing,—so much so that, as Prof. Schuh remarked upon another matter, "they have become 'massenhaft.'"

In all the different handbooks and treatises on ophthalmology we must expect to have many repetitions on the same subject, but at the same time we gain more fully the partic-

ular experience of each individual writer than in any other way; and the greater the experience the greater will be the gain to the reader and student. As in the case before us, Dr. Schweigger was for many years connected with the late v. Graefe as ophthalmoscopist in his large clinic in Berlin, and had with him an experience of great and lasting benefit. After the death of v. Graefe, he was appointed to fill his place as Professor of Ophthalmology in the University of Berlin, and surgeon in charge of the Ophthalmological Department in the Charité Hospital.

As a handbook is but a short manual of reference, it is not to be expected that we will find everything taken up and exhaustively treated. It is the purpose only of the author to call attention to the general matters in ophthalmology, and to enlarge upon the more important points, so as to give a good insight into the many different affections with sufficient information to enable the physician to diagnose the different forms of diseases and their treatment, as well as a foundation for pursuing his studies deeper and more thoroughly in this branch if so desired.

It is written in a short, concise, and very readable style, making it interesting and well calculated to prevent the reader from becoming wearied. Everything that the author deems necessary to explain is clearly and plainly elucidated in the briefest but most comprehensive manner, making it a valuable book for the busy practitioner. He excludes everything that would be of interest only to those who desire to make a special study of ophthalmology.

The translation is well done, the clear concise language of the original having been admirably preserved. It is, however, much to be regretted that he did not also act as English editor, and bring the contents up to the standard of the present day, by inserting in notes the many new and important points in relation to the pathology, treatment, and operations since the issue of the German work; for we find nothing new since 1874, and then only two references for that year. Really there are but ten references to new matters since the first edition in 1871.

The book is gotten up in most excellent style, with good paper and print, and is well worthy a place in the library of every physician, for reading and ready reference.

DISEASES OF THE NERVOUS SYSTEM. By SAMUEL WILKS, M.D., F.R.S. Philadelphia. Lindsay & Blakiston.

This is a well-written book of nearly 500 pages, containing little or nothing that is novel, but yet not lacking in the freshness which arises from the author having seen and studied at the bedside what he is talking about. It covers very fairly the field of nervous diseases, and we recommend it cordially as a guide to those who want to get a sufficient knowledge

of the subject for use in ordinary practice, but are unwilling or unable to give the time required by some of the more extensive works.

GLEANINGS FROM EXCHANGES.

UNUSUAL OCCURRENCE IN THE ATTEMPT TO REDUCE A DISLOCATED HUMERUS (*The Lancet*, July 6, 1878).—Mr. Thomas Smith reports the following very rare and interesting case, premising it with the remark that the gentleman who made the extension was not an athlete, nor was he supposed to possess any extraordinary physical force, neither had he put out his entire strength at the time of the accident: "A man, aged fifty-eight, was admitted to the hospital April 30, 1877. Eight weeks before admission he fell on his elbow, his hands being in his pockets at the time. As the result of this accident he suffered from paralysis of the radial nerve, and his arm lost its ordinary mobility, for which he was treated in various ways. On examination at the time of his admission he was found to have suffered a subcoracoid dislocation of the left humerus, which had escaped notice. On May 3, with the concurrence of the surgical staff of the hospital, he was put under the influence of ether, and after Mr. Smith had manipulated the limb so as to break down recent adhesions, an attempt was made to reduce the dislocation. The house-surgeon, sitting on the ground by the side of the patient, placed his left foot, covered only by a thin sock, in the axilla; a jack-towel was fastened by a clove hitch round the arm just above the elbow, the other end of the towel being passed behind the house-surgeon's shoulder, who also made extension by pulling from the patient's wrist. No other force was employed, and no assistance was given by bystanders. After extension had been maintained for a minute or so, the whole of the anterior fold of the axilla, integuments and muscles, gave way like 'rotten leather.' The cavity of the axilla was laid widely open, and pretty free hemorrhage took place. This was at once arrested by pressure, the patient was lifted on to the operating-table, the bleeding vessels were secured, the axilla was washed out with carbolic solution, the wound was partially closed, and drainage established by lint soaked in carbolic lotion.

"The pectoral muscles were found to have been almost completely torn, and the large vessels and nerves, with the head of the humerus, were laid bare, but not torn; the dislocation was reduced with ease. No immediate constitutional disturbance followed the accident. The patient took his food well, and his temperature was normal for four days; but as the discharge set in, his strength began to fail, and he died on the ninth day, from exhaustion.

"On post-mortem examination, diffuse sup-puration was found to exist in and around the axilla, and the parts about the upper and middle lobes of the right lung were in a state of consolidation. The heart was flabby; the liver large, pallid, and fatty; the kidneys normal; the spleen large, soft, and semi-fluid; vessels of the size of the tibials were rigid from calcareous degeneration. The muscles generally were paler, softer, and more flabby than normal. At the seat of injury nothing could be ascertained as to their condition as regards degeneracy, owing to the amount of sloughing that had taken place. No microscopical examination was made."

OPIMUM HABITUATION (*The Medical Record*, July 27, 1878).—Dr. J. B. Mattison, of Parrish Hall, Brooklyn, an institution exclusively devoted to the treatment of the victims of the opium-habit, details some cases coming under his notice, and thus summarizes the principles of treatment there adopted. Tonics and sedatives constitute the therapeutic reliance. The effect of preliminary *sedative* treatment in obviating the discomfort of the withdrawal of the opium is so satisfactory that we are most firmly convinced of its value.

Bromide of sodium is preferred for its pleasanter taste. It is given, largely diluted, three—occasionally four—times in the twenty-four hours. The plan of frequent and prolonged administration—thirty to sixty grains every half-hour, etc.—is not regarded with favor. The effect of the bromide being rather remote, this method incurs the risk of surcharging the system and entailing unpleasant if not serious results.

The efficacy of hot baths has been alluded to. Their value is beyond dispute.

For removal of the muscular debility we have found nothing equalling general faradization, twenty minutes sittings daily. The sense of exhilarating comfort resulting is very decided.

Strychnia is always employed with cod-liver oil, phosphorus, iron, digitalis, and quinine, as the case may demand; and out-of-door exercises, especially pedestrian, with varied social enjoyments, form valued adjuncts in treatment.

Our experience proves conclusively the efficacy of therapeutical resources in overcoming this formidable disorder, and warrants a favorable prognosis in all cases not complicated by organic disease.

THE ACTION OF SULPHURETTED WATERS IN REPRODUCING THE SYMPTOMS OF SYPHILIS WHEN LATENT (*The Practitioner*, July, 1878).—The explanation of the effects of sulphuretted waters in reproducing the symptoms of syphilis is stated by Güntz as follows: 1. A decomposition of albumen is produced by the water. 2. This decomposition is expressed clinically by a transitory chlorotic condition, owing to diminished size of the liver and spleen, and by increased excretion of urea.

3. This increase of urea is due to the action of the sulphuretted water in withdrawing oxygen, the effect of which is increased by coincident diminution of the activity of the heart and respiration. 4. The formation of new syphilitic products is effected at the cost of metamorphoses of albuminous substances, which by their decomposition produce new conditions, under which the syphilitic poison can act and form new combinations. 5. As long as the capacity of syphilitic contagion to attach itself to organic elements exists, so long can the poison remain in the body. Mercury acts in the same way as sulphuretted water, since it forms an albuminate in the body.

OFFICIAL LIST

OF CHANGES OF STATIONS AND DUTIES OF OFFICERS OF THE MEDICAL DEPARTMENT U.S. ARMY FROM AUGUST 25 TO SEPTEMBER 7, 1878.

KING, WILLIAM S., LIEUTENANT-COLONEL AND SURGEON.—Assigned to duty at Carlisle Barracks, Pa. S. O. 156, Department of the East, August 31, 1878. Granted leave of absence for six months, on account of sickness, with permission to apply for an extension, if necessary. S. O. 190, A. G. O., September 2, 1878.

SIMONS, J., LIEUTENANT-COLONEL AND SURGEON.—Assigned to duty at Fort McHenry, Md. S. O. 156, c. s., Department of the East.

HEAD, J. F., LIEUTENANT-COLONEL AND SURGEON.—Assigned to duty at Fort Independence, Mass. S. O. 156, c. s., Department of the East.

MCPARLIN, T. A., MAJOR AND SURGEON.—Assigned to duty at Fort Hamilton, N. Y. H. S. O. 156, c. s., Department of the East.

FRYER, B. E., MAJOR AND SURGEON.—Granted leave of absence for twenty-two days. S. O. 153, Department of the Missouri, August 26, 1878.

MCCLELLAN, E., MAJOR AND SURGEON.—When relieved by Assistant-Surgeon Gray, to proceed to Fort Lapwai, Idaho, and report for duty at that post. S. O. 104, Department of the Columbia, August 22, 1878.

NOTSON, W. M., MAJOR AND SURGEON.—When relieved by Assistant-Surgeon Koerber, to comply with S. O. 176, c. s., A. G. O. S. O. 176, Department of Dakota, August 20, 1878.

BROWN, H. E., CAPTAIN AND ASSISTANT-SURGEON.—Granted leave of absence for one month, with permission to leave the Department. S. O. 181, Department of Texas, August 26, 1878.

BARTHOLOMEW, J. H., CAPTAIN AND ASSISTANT-SURGEON.—Relieved from duty at Camp Harney, and assigned to duty at Fort Stevens, Oregon. S. O. 104, c. s., Department of the Columbia.

KOERPER, E. A., CAPTAIN AND ASSISTANT-SURGEON.—Assigned to duty at Fort McKinney, Wyo. T., relieving Major Notson. S. O. 77, c. s., Department of Dakota.

CLEARY, P. J. A., CAPTAIN AND ASSISTANT-SURGEON.—Leave of absence extended one month. S. O. 65, Division of the Missouri, August 26, 1878.

STEINMETZ, WM. R., CAPTAIN AND ASSISTANT-SURGEON.—Granted leave of absence for one year. S. O. 186, A. G. O., August 28, 1878.

GEAY, W. W., FIRST LIEUTENANT AND ASSISTANT-SURGEON.—Assigned to temporary duty at Fort Vancouver, W. T. S. O. 104, c. s., Department of the Columbia.

BRECHMIN, L., FIRST LIEUTENANT AND ASSISTANT-SURGEON.—Assigned to duty as Post-Surgeon at the new post to be established near Camp "J. G. Sturgis" (Bear Butte, D. T.) by Major Lazelle, First Infantry. S. O. 102, Department of Dakota, August 27, 1878.

CAMPBELL, A. B., CAPTAIN AND ASSISTANT-SURGEON.—Died at Chicago, Ill., on September 1, 1878.

WYETH'S DIALYSED IRON.

(FERRUM DIALYSATUM.)

A Pure Neutral Solution of Peroxide of Iron in the Colloid Form. The Result of Endosmosis and Diffusion with Distilled Water.

PREPARED SOLELY BY

JOHN WYETH & BRO.,
PHILADELPHIA.

This article possesses great advantages over every other ferruginous preparation heretofore introduced, as it is a solution of Iron in as nearly as possible the form in which it exists in the blood. It is a preparation of invariable strength and purity, obtained by a process of dialysation, the Iron being separated from its combinations by endosmosis, according to the law of diffusion of liquids. It has no styptic taste, does not blacken the teeth, disturb the stomach, or constipate the bowels.

It affords, therefore, the *very best* mode of administering

IRON

in cases where the use of this remedy is indicated.

The advantages claimed for this form of Iron are due to the absence of free acid, which is dependent upon the perfect dialysation of the solution. The samples of German, French, and American Liquor Ferri Oxidi Dialys., which we have examined, give acid reaction to test paper. If the dialysation is continued sufficiently long, it should be tasteless and neutral.

Our Dialysed Iron is not a saline compound, and is easily distinguished from Salts of Iron by not giving rise to a blood-red color on the addition of an Alkaline Sulpho-Cyanide, or a blue precipitate with Ferro-Cyanide of Potassium. It does not become cloudy when boiled. When agitated with one part of Alcohol and two parts of Ether (fortior), the Ether layer is not made yellow.

Physicians and Apothecaries will appreciate how important is the fact that, as an antidote for poisoning by Arsenic, Dialysed Iron is quite as efficient as the Hydrated Sesquioxide (hitherto the best remedy known in such cases), and has the great advantage of being always ready for immediate use. It will now doubtless be found in every drug store to supply such an emergency.

FULL DIRECTIONS ACCOMPANY EACH BOTTLE.

Physicians will find our DIALYSED IRON in all the leading drug stores in the United States and Canada.

It is put up in bottles retailing for ONE DOLLAR, containing sufficient for four months' treatment. Large size is intended for hospitals and dispensing. Retail at \$1.50. Price Lists, etc., etc., sent on application.

JOHN WYETH & BRO.

WYETH'S COMPRESSED TABLETS OF CHLORATE OF POTASH.

For Hoarseness, Bronchial Irritation, Sore Throat, Diphtheria, Croup, etc., etc.

Chlorate of Potash is a remedy of acknowledged value in cases of Diphtheritic Sore Throat, and in Inflammation of the Mouth and Throat, induced by a depressed state of the system. In these instances, as in the milder forms of Croup, it has, besides its depurative and detergent effects, a solvent action on the deposits characteristic of those troublesome and dangerous affections. It relieves Hoarseness; and in many cases of Fetid Breath from disordered secretions, it proves an efficient corrective. Its virtues in simple Angina, or ordinary Sore Throat, are recognized by many of the most eminent physicians.

As the taste of this article is not disagreeable, we have prepared it in the form of Compressed Tablets, thus giving the patient the full benefit of its action, undiluted with Sugar, Gum, or other vehicles, which would not only prevent its effects, but which sometimes themselves offend the stomach.

The Lozenges usually contain about twenty five (25) grains of gum and sugar, with two grains of the Chlorate of Potash, while each of these Pills contains simply five grains of the Chlorate, all of which, dissolved in the saliva, acts on the affected mucous membranes.

If allowed to dissolve in the mouth, the topical effect is much more efficient than a saturated solution, as while the solution is but temporary, the tablet really acts as a continuous gargle.

FOR SORE THROAT, HOARSENESS.

DIRECTIONS.—Adults should take one every hour or two until relieved, allowing it to dissolve slowly in the mouth. Children, half of one as often.

For Offensive Breath, no remedy will give more certain relief; use one two or three times a day.

For Diphtheria, Croup, and the more serious ailments, the physician should direct.

P. SCHERER & CO.,

Successors to JOHN SATTIG. (Established 1841.)

Nos. 72 and 74 PARK PLACE, NEW YORK, AGENTS FOR

LÖBEL SCHOTTLÄNDER, Carlsbad, Bohemia,

RECOMMEND TO THE PROFESSION AND TRADE ESPECIALLY,

CARLSBADER SPRING WATER,
AND OTHER PRODUCTS OF THE SPRINGS.

According to our latest contracts with the proprietors of the different Carlsbad Springs, we get fresh shipments with every steamer, so as to be able to supply this water in a quality not to be had formerly. Carlsbad Waters, and especially the **SPRUDEL, SCHLOSS (Castle), THERESIA, MUEHL (Mill) SPRING**, are to-day relied on by the Medical Profession in Germany as the surest remedy in Diabetes, Diabetes Mellitus, and Diseases of the Kidneys. They are likewise used as a specific almost for **Reducing Obesity**. They are a mixed alkaline, saline water, of reliable healing quality in Liver and Bowel Complaints, and Disease of the Prostate Gland. Sold, **ALWAYS FRESH**, only by us. For price and other information, please send for Pamphlet on Mineral Waters. Special quotations on quantities.

CARLSBAD SPRUDEL SALT may be used either to assist the effects of the Carlsbad Water or as a remedy by itself. In the latter case the dose is one or two teas. spoons, either in a dry state or dissolved in water. It dissolves rapidly and completely in warm water, and in this form a curative effect may be expected with more certainty. It is advisable to use water at a temperature of 30° to 40° R. This, however, does not exclude its being taken with cold water. The best time for taking the salt is in the morning, shortly before breakfast, fasting.

CARLSBAD SPRUDEL SOAP, for Psoriasis, Eczema, and Chronic Dermatoses, always on hand.
CARLSBAD SPRUDEL LOZENGES, representing in a light degree the qualities of the water, always on hand, **FRESH IMPORTED.**

Keep continually fresh on hand Aachner, Aix-la-Chapelle, Apollinaris, Birmensdorfer, Carlsbader, Emser Kraenchen, Friedrichshaller, Gieshuebler, Haller Iodine, Homburger Elizabeth, Hungarian Bitterwasser, Kissinger Ragoczy, Marienbader Kreuzbrunnen, Ofener Ragoczy, Ofener Franz Joseph, Pullna Bitterwater, Pyrmont Steelwater, Roisdorfer, Schwalbacher Steelwater, Selters, Vichy, Wildungen Otto and Helenen, and many more. Send for Catalogue and Prices.

Agent in Philadelphia, **B. G. ELLINGER, 123 S. Front St.****F. ALFRED REICHARDT,**

Pharmacist and Importer, 404 Fourth Avenue, New York,

SOLE AGENT FOR**GARNIER, LAMOUREUX & CIE.,**

13 Rue St.-Honore, Paris,

the oldest and largest house in the world for

SUGAR-COATED PILLS and GRANULES.

We use nothing but pure Sugar for coating, free of Ialcum, Terra Alba, Gypsum, etc. Dragées of Crystallized Valerianate of Ammonia, Ferri Ponati, Monobromide of Camphor, Croton Chloral, Meglin, Dia cordinum, Phosphorus, etc., etc. Send for circular and price list.

DR. FRIEDRICH WITTE, Rostock, Germany,**CHEMICAL WORKS.****THE STRONGEST PEPSIN MADE,—****PEPSINUM PLANE SOLUBILE.**

One grain will dissolve 100 grains of Albumen, Pylalin. One grain will dissolve 100 grains of Starch or Starchy Food, Pancreatin, and Dry Extract of Rennet. Send for circular and prices.

MATHIAS ROZSNYAY,

Arad, Hungary.

TASTELESS QUININE

(Neutral Tannate), and

TASTELESS QUININE LOZENGES.

Absolutely free from bitter taste.

Recommended by the

ACADEMY OF MEDICINE IN PARIS.

Send for circular and price list.

FRANCIS J. GEIS, BREWER AND MALTSTER,

Dobbs Ferry, New York.

EXTRACT OF MALT.

The best in the world. I guarantee that this extract contains absolutely nothing but the soluble parts of the best Canada Barley Malt and Distilled Water. Contains more Albumen and Phosphoric Salts than any other, and challenges analysis. Retailed in 1-lb. stone jars at 50 cents.

PROPRIETOR OF**FLUID EXTRACT OF DAMIANA.**

(Late Helmick, or Helmick & Co., Washington.)

Made from the genuine Damiana leaf imported by myself, pronounced the best and most reliable of any in the market by Dr. J. J. Caldwell, of Baltimore; Prof. R. Hutchinson, Providence; Dr. Noah Sanborn, Bayonne, N. J.; Dr. G. Buckingham Smith, New York, and many more. Send for circular and prices. Sample gratis to physicians.

WARBURG'S TINCTURE.

Introduced by me in America. This remedy is now largely used by the best practitioners in the United States in cases where quinine fails to act. I guarantee this to contain every ingredient prescribed in the original formula. Send for circular and price list.

NITRITE OF AMYL PEARLS,

Made solely from

DR. E. R. SQUIBB'S PURE NITRITE OF AMYL.

Beware of imitations. They are the only ones correctly dosed, containing 2, 3, 5, 8, 10 drops, or made larger to order if desired. Highly recommended for Asthma, Epilepsy, Angina Pectoris, and SEA-SICKNESS. Send for circular and prices.

IODOFORM CRAYONS, ERGOT CRAYONS.

Brush-holding Rubber Stopples.

Elixir of Red Peruvian Bark, the most effective and best-tasting. Apol Capsules. Pure Etheral Cubebs Extract in Capsules. Fluid Extract of Rhubarb and Dandelion. Yerba Santa, Fucus Vesiculosus.

NITRITE OF AMYL PEARLS,

With Combination Box and Crusher, made so strong that they will not explode in hot weather. A certain remedy against SEA-SICKNESS. Order pamphlet, "Antidote against Chloroform Poisoning," *New York Medical Record* of July 20, 1878, article by Dr. Burrell.

These pearls were formerly blown of glass so thin that they could be easily crushed between the fingers without wounding them. The great objection, however, arose from the fact that the Nitrite of Amyl, being very volatile, became so much expanded by the slightest rise in the temperature that often 90 per cent. of the manufactured pearl would break.

"This latest difficulty is now successfully overcome by making them of glass so thick and strong that the Nitrite of Amyl cannot burst it, and providing an instrument for breaking the pearls with each box of one dozen pearls. This instrument, which is secured by letters patent, consists of two boxes, one fitting loosely into the other; between the two the pearl is laid in a little cotton."

Agent for the above, **B. G. ELLINGER, 123 S. Front St., Philadelphia.**

To the Medical Profession.

LACTOPEPTINE,

The most important remedial agent ever presented to the Profession for Indigestion, Dyspepsia, Vomiting in Pregnancy, Cholera Infantum, Constipation, and all Diseases arising from imperfect nutrition, containing the five active agents of digestion, viz.: Pepsin, Pancreatine, Diastase, or Veg. Ptyalin, Lactic, and Hydrochloric Acids, in combination with Sugar of Milk.

FORMULA OF LACTOPEPTINE.

Sugar of Milk	40 ounces.	Veg. Ptyalin or Diastase	4 drachms.
Pepsin	8 "	Lactic Acid	5 fluid drachms.
Pancreatine	6 "	Hydrochloric Acid	5 "

LACTOPEPTINE owes its great success solely to the Medical Profession, and is sold almost entirely by Physicians' Prescriptions. Its almost universal adoption by the profession is the strongest guarantee we can give that its therapeutic value has been most thoroughly established.

The undersigned, having tested LACTOPEPTINE, recommend it to the profession.

ALFRED L. LOOMIS, M.D.,
Professor of Pathology and Practice of Medicine, University of the City of New York.

F. LE ROY SATTERLEE, M.D., Ph.D.,
Professor of Chem., Mat. Med., and Therap. in the N. Y. College of Dent.; Prof. of Chem. and Hygiene in the Am. Vet. College, etc., etc.

LEWIS A. SAYRE, M.D.,
Professor of Orthopedic Surgery and Clinical Surgery, Bellevue Hospital Medical College.

SAMUEL R. PERCY, M.D.,
Professor Materia Medica, New York Medical College.

A. VAN DEVEER, M.D.,
Albany, N. Y., June 8, 1878.
Prof. of the Prin. and Prac. of Surg., Albany Med. Col.; Surg. Albany and St. Peter's Hospitals.

"I have given LACTOPEPTINE a good, thorough trial, and have been greatly pleased with the excellent results that have followed its administration."

JOHN H. PACKARD, M.D.,
Philadelphia, Pa., May 30, 1878.
Pres't Pa. Co. Obst. Society; Surg. Episcopal and Women's Hospitals.

"I have found great satisfaction in the use of LACTOPEPTINE, and have ordered it frequently in cases of Dyspepsia, especially where there is want of tone and defective secretion."

JAMES AITKEN MEIGS, M.D.,
Philadelphia, Pa., June 20, 1878.
Prof. of the Institutes of Med. and Med. Juris., Jeff. Medical College; Phys. to Penn. Hos.

"I have used LACTOPEPTINE with very good effect in a number of cases of Dyspepsia."

W. W. DAWSON, M.D.,
Cincinnati, Ohio, June 21, 1878.
Prof. Prin. and Prac. Surg. Med. Col. of Ohio; Surg. to Good Samaritan Hospital.

"I have used LACTOPEPTINE with great advantage in cases of feeble digestion."

ALBERT F. A. KING, M.D.,
Washington, D. C., June 19, 1878.
Prof. of Obstetrics, University of Vermont.

"I have used LACTOPEPTINE both in hospital and private practice, and have found it to answer fully the purpose for which it is recommended. As an immediate aid to the digestive function, I know of no remedy which acts more directly."

D. W. YANDELL, M.D.,
Louisville, Ky., March 7, 1878.
Prof. of the Science and Art of Surg. and Clinical Surg., University of Louisville.

"I have made much use of LACTOPEPTINE, and take great pleasure in stating that it has rarely disappointed me. I shall, of course, continue to prescribe it."

ROBT. BATTEY, M.D.,
Rome, Ga., June 7, 1878.
Emeritus Prof. of Obstetrics, Atlanta Med. College, and ex-Pres. Med. Association of Ga.

"I have used LACTOPEPTINE in a case of Dyspepsia with satisfaction. I think well of it."

CLAUDE H. MASTIN, M.D., LL.D.,
Mobile, Ala., June 8, 1878.

"I consider LACTOPEPTINE the very best preparation of the kind which I have ever employed, and for patients with feeble digestion I know of nothing which is equal to it."

Prof. H. C. BARTLETT, Ph.D., F.C.S.,
London, England,
February 22, 1876.

"I find the preparation of LACTOPEPTINE contains within itself all the principles required to promote a healthy digestion."

PRICE LIST.

LACTOPEPTINE (in oz. bottles).....	per oz.	\$1.00
" (in 4 oz. ").....	per doz.	12.00
" (in 1/2 lb. ").....	per lb.	12.00

"We also prepare the various Elixirs and Syrups in combination with Lactopeptine.

THE NEW YORK PHARMACAL ASSOCIATION,

2 & 3 COLLEGE PLACE, NEW YORK.

P. O. BOX 1574.

TROMMER'S EXTRACT OF MALT.

The rapidly increasing demand for our IMPROVED EXTRACT OF MALT, during the four years that it has been manufactured and offered to the medical profession in America, justifies the belief that in its production here we are meeting a generally felt want.

Long experience in manufacturing Malt Extract has enabled us to completely overcome the many difficulties attending its manufacture in large quantity; and we positively assure the profession that our Extract of Malt is not only perfectly pure and reliable, but that it will keep for years, in any climate, without fermenting or moulding, and that its flavor actually improves by age. Our Extract is guaranteed to equal in every respect the best German make, while, by avoiding the expenses of importation, it is afforded at less than half the price of the foreign article.

The Malt from which it is made is obtained by carefully malting the very best quality of selected Toronto (Canada) Barley. The Extract is prepared by an *improved process*, which prevents injury to its properties or flavor by excess of heat. **It represents the soluble constituents of Malt and Hops**, viz., MALT SUGAR, DEXTRINE, DIASTASE, RESIN and BITTER OF HOPS, PHOSPHATES OF LIME and MAGNESIA, and ALKALINE SALTS.

Attention is invited to the following analysis of this Extract, as given by S. H. Douglas, Professor of Chemistry, University of Michigan, Ann Arbor.

TROMMER EXTRACT OF MALT CO.—I enclose herewith my analysis of your Extract of Malt:

Malt Sugar, 46.1; Dextrine, Hop-bitter, Extractive Matter, 23.6; Albuminous Matter (Diastase), 2.469; Ash—Phosphates, 1.712; Alkalies, .377; Water, 25.7; Total, 99.958.

In comparing the above analysis with that of the Extract of Malt of the German Pharmacopœia, as given by Hager, that has been so generally received by the profession, I find it to substantially agree with that article.

Yours truly,

SILAS H. DOUGLAS,

Professor of Analytical and Applied Chemistry.

This invaluable preparation is highly recommended by the medical profession as a most effective therapeutic agent for the restoration of delicate and exhausted constitutions. It is very nutritious, being rich in both muscle- and fat-producing materials.

The very large proportion of *Diastase* renders it most effective in those forms of disease originating in *imperfect digestion of the starchy elements* of food.

A single dose of the Improved Trommer's Extract of Malt contains a larger quantity of the active properties of malt than a pint of the best ale or porter; and, not having undergone fermentation, is absolutely free from alcohol and carbonic acid.

The dose for adults is from a dessertspoonful to a tablespoonful three times daily. It is best taken after meals, pure or mixed with a glass of milk, or in water, wine, or any kind of spirituous liquor. Each bottle contains 1½ lbs. of the Extract.

Our preparations of Malt are for sale by druggists generally throughout the United States and Canada, at the following prices.

EXTRACT OF MALT, <i>With Hops (Plain)</i>	\$1.00
“ “ “ <i>Phosphate of Iron (Ferrated)</i>	1.00
“ “ “ <i>Cod Liver Oil</i>	1.00
“ “ “ <i>Cod Liver Oil and Iodide of Iron</i>	1.00
“ “ “ <i>Cod Liver Oil and Phosphorus</i>	1.00
“ “ “ <i>Hypophosphites</i>	1.50
“ “ “ <i>Iodides</i>	1.50
“ “ “ <i>Alteratives</i>	1.50
“ “ “ <i>Citrate of Iron and Quinia</i>	1.50
“ “ “ <i>Pepsin</i>	1.50

MANUFACTURED BY
TROMMER EXTRACT OF MALT CO.,
FREMONT, OHIO.

SUNNYSIDE.

A PRIVATE MEDICAL HOME FOR NERVOUS INVALIDS.

Washington Heights, New York.

This Home, unsurpassed for its beauty of location and surroundings, and with handsomely furnished apartments, receives a limited number of cases, of either sex, of *Mental and Nervous Diseases*, *Dipsomania*, and *Opium-Habit*, and offers to such patients all the attentions, comforts, and attractiveness of a home, together with medical treatment. Applications may be made either by mail or personally at 152d St., bet. 10th Ave. and Boulevard. Opium-Habit and Dipsomania cured without suffering. Billiards, Driving, Rowing, and Horseback Exercise.

Twelve trains daily from Hudson River Depot, 30th St. and 10th Avenue. Dr. EDWARD C. MANN, M.D., Medical Superintendent.

QUASSIA CUPS.

"SIMARUBA EXCELSA."

We would respectfully call the attention of the medical profession to the real merits of these medicinal goblets.

It is an established fact that, for a healthy tonic, it has not an equal in the world. For fever and ague it is a quick and sure remedy.

A cure for general debility and loss of appetite.

We manufacture the cups in three sizes. Full directions with each cup.

For the genuine, observe the name on the label.

The wood is imported by us direct from the West Indies, and thus we can warrant the quality.

Send for circular and price-list, or 50 cents for sample cup.

Address all communications to

THE

Peru Medicinal Goblet Manufactory,

H. W. McINTIRE,

Manufacturers' Agent,

SPRINGFIELD, MASS.

SHEDDEN'S FOODS FOR THE DYS-

PEPTIC AND DIABETIC. Diabetic Flour, free from Starch, 10 lb. cases, \$5. Prepared Gluten Flour, 10 lb. cases, \$3. Depot, 1311 Broadway, corner 34th St., New York City.

The Penn Mutual Life Insurance Company of Philadelphia.

ASSETS ACCUMULATED, \$6,280,723.46.

The Penn is a purely mutual company. All of its surplus premiums are returned to the members every year. Dividends may be applied to reduce the premium or increase insurance as desired.

SAMUEL C. HUEY, President.

H. S. STEPHENS, Vice-President. H. AUSTIE, Secretary.

EDWARD A. PAGE, M.D., Medical Director.

EDWARD HARTSHORNE, M.D., Consulting Med. Examiner.

WOMAN'S MEDICAL COLLEGE

OF THE

NEW YORK INFIRMARY,

128 Second Avenue, Cor. 8th St., New York.

Regular Session of 1878-79 will open October 1st. A thorough course of three years' instruction by Lectures, Clinics, and practical work in Laboratory, Drug-Room, and different Wards of the Infirmary. For Catalogues, etc., address the Secretary of the College;

MERCY N. BAKER, M.D.,
128 Second Avenue, New York City.

CLEVELAND MEDICAL COLLEGE.

The Thirty-sixth Session will commence on the 18th of Sept., 1878, and continue twenty-four weeks. The Summer Session commences on the first Monday in April.

Faculty.—J. L. Cassels, M.D., LL.D., Chemistry, Emeritus; Proctor Thayer, M.D., Surgery; H. K. Cushing, M.D., Gynecology; John Bennett, M.D., Dean, Practice; Jacob Laisy, M.D., Osteology; John E. Darby, M.D., Therapeutics; E. W. Morley, M.D., Ph.D., Chemistry; Isaac N. Himes, M.D., Physiology; N. C. Scott, M.D., Ophthalmology; B. W. Holliday, M.D., Anatomy; Lewis Buffett, M.D., Oral Surgery; H. H. Powell, M.D., Obstetrics; W. W. Holliday, M.D., Demonstrator.

Fees.—Lecture tickets, including Matriculation, \$50.

For Catalogue, address

HUNTER H. POWELL, M.D., Sec.,
323 Euclid Avenue.

Woman's Medical College

OF

PENNSYLVANIA.

The Twenty-Ninth Winter Session will open on Thursday, October 3, 1878, in the commodious new college building.

Clinical instruction is given in the Woman's Hospital, Pennsylvania, Wills, Philadelphia, and Orthopaedic Hospitals.

Spring Course of Lectures, Practical Demonstrations, and Winter Quizzes are free (except for expense of material) to all matriculants of the year.

Address RACHEL L. BODLEY, A.M., Dean,
North College Ave. and 21st St., Philadelphia, Pa.

Pennsylvania College of Dental Surgery.

Twenty-Third Annual Session, 1878-79.

The regular Winter Session for 1878-79 will commence on Tuesday, October 1, 1878, and continue five months, in the new and commodious college building, on Twelfth Street, below Arch, to which the College has just been removed. The size and equipment of the rooms in this building will enable the Faculty to offer the Dental student the advantages accruing from lecture-rooms with high ceilings and comfortable seatings for over two hundred (200) students, clinic-rooms with such an abundance of light that the most delicate operations can be performed in any part of the room, well-furnished laboratories,—mechanical and chemical, and a large and well-lighted and well-ventilated dissecting-room. For further information, address

C. N. PIERCE, Dean,
1617 Green Street, Phila.

Toussaint's Concentrated Extract of Malt.

Prepared from the best Canada Barley Malt. Evaporated in a vacuum apparatus at a temperature below 120° F. The attention of the Medical Profession and the trade is called to my preparations of **Extract of Malt** (plain), as also to the compounds.

Send for samples, price list, and circulars, giving full information.

FRED TOUSSAINT, Chemist, New York.

Office, 130½ Essex St. Laboratory, 134 Essex St.

Wholesale Depot at ASCHENBACH & MILLER,
Third and Callowhill Streets, Philadelphia.

VACCINE VIRUS.

In order to meet properly the continued demand upon us for **Animal Virus**, we have established stables for its propagation upon carefully-selected heifers. The lymph used is of the well-known "Beaugency" stock, imported by ourselves expressly for this purpose. The result of several years' experience in supplying this lymph leads us to believe that its excellence is unsurpassed. The establishment is under the care of a competent physician, who will spare no pains to produce a perfectly **reliable** and **pure** article, which we are prepared to furnish **fresh, daily**.

We can also furnish, to those who prefer it, **Humanized Virus**, from healthy children, procured for us by physicians of undoubted reliability.

All our Virus is put up in strong, *air-tight, sealed packages*, for safe conveyance by mail or express, and will be sent (postpaid if by mail) upon the following terms:

From the Heifer,	10 Large Ivory Points, well charged on both sides		\$1.50
	5 Large Ivory Points, " " " "		.80
	Large Ivory Points, less than 5, well charged on both sides,	each	.25
	1 Crust, new method, in air-tight Glass Capsule		2.00

From Healthy Infants,	10 Small Ivory Points		1.50
	1 Crust from Unruptured Vesicles		2.00

Directions for vaccinating with either form of Virus, derived from methods successfully employed, will be furnished with Virus if requested.

We will warrant every package of Points and every Crust, giving a fresh supply in case of failure reported within fifteen days for Points, thirty days for Human, and ninety days for Kine Crusts. We can usually furnish Crusts one remove from the heifer if preferred.

On account of their unreliability, we have hitherto furnished the usual form of Kine Crust unwillingly. Under our new method of taking and preserving them, however, after careful tests and an experience extending over several months, the results attained have been so satisfactory that we now offer them as not less active and reliable than other forms of Virus, while less liable to become inert with lapse of time. We now offer them on very favorable terms, and recommend them for transmission to a distance and in all cases where it is desired to preserve Virus for some weeks or months, or to keep a supply at hand for emergencies.

We also furnish **Uncharged Ivory Points**, for physicians' use, at the following rates:

Small	per 100, 25 cents; per 1000, \$2.00
Large	per 100, 50 cents; per 1000, 4.00

Orders by mail or telegraph answered by return train.

Liberal discounts upon large supplies for Cities, Towns, and Institutions.



Scarifying Vaccinator. Steel, Nickel-plated. (See cut.) Each 25 cents.

New Illustrated Catalogue of Surgical Instruments, postpaid, on request.

CODMAN & SHURTLEFF,

MAKERS AND IMPORTERS OF

SUPERIOR SURGICAL INSTRUMENTS,

13 and 15 Tremont Street, Boston.

LIQUEFIED NITROUS OXIDE.

We are prepared to supply Liquefied Nitrous Oxide, of our own manufacture, *warranted pure*, in four sizes of cylinders, viz., 100, 700, 1000, and 1400 gallons.

Surgeons' Cases of stout morocco, neatly finished and lined with velvet, containing Cylinder, Tubing, Gas Bag, and Inhaler.

Cylinders refilled with Gas. A descriptive circular sent on application.

SAMUEL S. WHITE,
CHESTNUT STREET, COR. TWELFTH, PHILADELPHIA.

THE SPERMATIC TRUSS.

A most efficient, safe, and simple appliance for the treatment of Spermatorrhœa and Subduing Inordinate Sexual Desire. Entirely mechanical; operates by taking advantage of physical action.

THE RESULT OF YEARS OF PRACTICAL EXPERIENCE.

COMMENDED BY PHYSICIANS AND MEDICAL JOURNALS.

FAR SUPERIOR TO ANYTHING HERETOFORE USED FOR THAT PURPOSE.

WE EXCEPT NO MEDICATION OR APPLIANCE WHATEVER.

Price to physicians, \$4.00. For Truss, or particular description, address

COOPER TRUSS CO.,

P. O. Lock Box 55, PITTSBURGH, PA.

AN IMPORTANT WORK NOW READY.

STRICTURE OF THE MALE URETHRA: ITS RADICAL CURE.

BY FESSENDEN N. OTIS, M.D.,

Professor of Genito-Urinary Diseases in the College of Physicians and Surgeons, Surgeon to Charity Hospital, and President of the Medical Board, New York City, etc., etc.

OCTAVO. 358 PAGES. VERY FULLY ILLUSTRATED. \$3.00.

To the accomplished author is the medical profession mainly indebted for the most rational and effective method of treating strictures of the urethra; and now in this volume he not only presents the conclusions of nearly two decades of close study and observation, but he traces lucidly and in an interesting manner the gradual progress of his subject up to its present development. The causes, symptoms, complications, and results of stricture are indicated with great force; the old and—it must be said—*inefficient* modes of treatment are thoroughly analyzed; and the measures, which in nearly all cases give assurance of radical cure, are so ably presented and sustained by statistical proofs that few will be bold enough to dispute the claims urged for them. The tables which complete the work must carry conviction to every unbiased mind, and the work as a whole cannot fail to become a standard authority upon the subject of which it treats.

G. P. PUTNAM'S SONS, Publishers, 182 Fifth Avenue, New York.

BOSTON UNIVERSITY SCHOOL OF MEDICINE. OPEN TO BOTH SEXES.

THE lectures of the sixth year will begin October 9, 1878, and continue till June, 1879. This Medical School was one of the first in this country to require preliminary examinations, and to furnish a three years' graded course. To secure still more thorough results an optional *four years' course* is established, by which the graduate attains an additional degree in medicine as evidence of study. Address

I. T. TALBOT, M.D., Dean,
66 Marlborough St., Boston.

Every Man
HIS OWN
Printer!  **THE \$3 Press** Prints In-
etc. (Self-inked &c. &c. larger sizes
for business, pleasure, young and old
Catalogue of Presses, Types, Etc.,
for 2 stamps—KELLY & Co.
Meriden, Conn.

NOW READY.

University Hospital Clinics.

BEING CLINICAL LECTURES

On the Rational Treatment of Typhoid Fever.
By Prof. WM. PEPPER, M.D.

On Abortion: Its Symptoms and Treatment.
By Prof. R. A. F. FENKES, M.D.

On Sexual Exhaustion.
By Prof. HORATIO C. WOOD, Jr., M.D.

On the Operation for Laceration of the Female Perineum.

8vo. Paper Cover. 25 Cents.

*For sale by Medical Booksellers generally, or will be sent, postpaid, upon receipt of price by

J. B. LIPPINCOTT & CO., Publishers,
715 and 717 Market St., Philadelphia.

THE BUFFALO LITHIA WATER,

For Gout, Paralysis, and Affections Peculiar to Women.



of inveterate chronic gout has afforded much satisfaction to us, free diuresis being provoked and followed by relief of symptoms."

GOUT.

[DR. HORATIO C. WOOD, of Philadelphia, Professor of Materia Medica in the University of Pennsylvania. From editorial in the Medical Times of July 20, 1878.]

"Buffalo Lithia Springs.

"The value of alkalis in disease has long been known, and the wide reputation of the springs of Vichy bears testimony to the superiority of the natural waters over the simple alkalis. Of late years lithia has been asserted to have especial value in chronic gout over and above that of other alkalis. Some considerable experience has indicated that this assertion is well founded; and several years since we looked for a native lithia water which should be cheap and efficient. The product of the Buffalo Springs, of Mecklenburg county, Va., was finally brought to our notice by a Baltimore physician who had been relieved by its use of some very troublesome and alarming symptoms believed to be due to an inherited *gouty diathesis*. Trial in one or two cases

RHEUMATIC GOUT.

[DR. HUNTER MCGUIRE, Professor of Surgery in the Richmond (Va.) Medical College.]

"I have used the Buffalo Lithia Waters in cases of old rheumatic gout which had resisted the ordinary remedies, with wonderfully good results. I have myself suffered a great deal lately from this malady. The Lithia Water certainly does me more good than anything else I have used."

PARALYSIS, DYSPEPSIA, GASTRALGIA, Etc.

Case of Miss F—, stated by Dr. William B. Towles, Demonstrator of Anatomy in the Medical Department of the University of Virginia, and Resident Physician at these Springs.

"BUFFALO LITHIA SPRINGS, August 5, 1878.

"Miss F—, aged thirty, was examined June the 15th, 1878, and the following history of her case elicited: For several years she had experienced repeated attacks of intermittent fever, which had produced great debility, attended by frequently recurring paroxysms of functional disease of the heart; by dyspepsia and gastralgia so intense that vegetable food had been entirely excluded from her diet. There had also been paralysis of the lower limbs to such an extent that she had been forced to have recourse to crutches.

"The examination revealed marked œdema of both upper and lower limbs, and of the face; marked anæmia, mental depression, and gastric pains; considerable lateral curvature of the spine, and entire inability to maintain the erect position without crutches. The urine was scanty, high-colored, loaded with mucus, and depositing the urates copiously. Directions were given her for the use of the waters, and her diet regulated. Immediate improvement was evidenced, which has continued up to the present time. She has now thrown aside her crutches, walking readily without them; is able to eat with impunity any article of diet, and would be pronounced well but for slight spinal curvature.

"WILLIAM B. TOWLES, M.D."

COLD (LITHIA) WATER vs. HOT WATER FOR PARALYSIS AND AFFECTIONS OF THE KIDNEYS AND BLADDER.

THE BUFFALO LITHIA SPRINGS GIVE RELIEF AFTER A FAILURE OF THE CELEBRATED HOT SPRINGS OF ARKANSAS.

Case of Dr. P. S. Jones, a retired physician of Memphis, Tennessee.

[Extract from a letter of Dr. JONES, dated Memphis, August 3, 1878.]

"Awaking from sleep in a current of cold air, having previously enjoyed good health, I found myself affected with severe pain in the loins, which continued for seventy-two hours, the pain extending on one side along the great sciatic nerve to the extremity of the limb. With the subsidence of these symptoms I found that there was paralysis of the limb. After three years of suffering, which was attended by painful contraction of the muscles, frequently so great as to necessitate the use of opiates to produce sleep, an affection of the kidneys and bladder supervened; symptoms, frequent and urgent demand for voiding the urine, perhaps as often as every twenty or thirty minutes, which was highly colored and charged with blood, its passage being attended with great burning and spasm of the neck of the bladder, unfitting me for business. I was treated by several eminent physicians and surgeons, and made several visits to the celebrated Hot Springs of Arkansas, fully testing these waters; but all without material benefit. I have now been using for some weeks the waters of the Buffalo Lithia Springs, Va., with astonishing results. I am so far relieved of all my distressing symptoms, above stated, that I suffer no inconvenience or concern about my health. I sleep well, eat heartily, walk readily without a cane, attend regularly to my business affairs, and have no doubt of a speedy, perfect, and permanent cure in all respects from the further use of the water.

Respectfully,

P. S. JONES, M.D."

BUFFALO LITHIA WATER.

PECULIAR AFFECTIONS OF WOMEN.

[Extract from letter of Dr. WM. T. HOWARD, of Baltimore, Professor of Diseases of Women and Children in the University of Maryland, dated August 1, 1873.]

Dr. Howard compares these Waters with those of the celebrated Greenbrier White Sulphur, West Virginia, and adds the following:

"Indeed, in a certain class of cases, it is much superior to the latter. I allude to the abiding debility attendant upon the tardy convalescence from grave acute diseases; and more especially to the cachexia and sequels incident to malarious fevers in all their grades and varieties, to certain forms of atonic dyspepsia, and ALL THE AFFECTIONS PECULIAR TO WOMEN that are remediable at all by mineral waters. In short, were I called upon to state from what mineral waters I have seen the greatest and most unmistakable amount of good accrue in the largest number of cases in a general way, I would unhesitatingly say the Buffalo Springs, in Necklenburg county, Virginia."

[Extract from letter of Dr. HARVEY L. BYRD, of Baltimore, late Professor of Diseases of Women and Children in the College of Physicians and Surgeons of Maryland, dated Baltimore, September 21, 1877.]

"During my stay at the Buffalo Lithia Springs, as resident physician, for the season just closed, I had ample opportunities for observing the action of the waters of the several Springs on different constitutions and the two sexes, and I have no hesitation in expressing the opinion that, in the large majority of chronic FEMALE DISEASES, and especially those connected with the Uro-Genital Organs, they have no equal as a medicinal agent in the whole range of Therapeutic application. In Leucorrhœa, Amenorrhœa, Dysmenorrhœa, and Cystorrhœa their action may be regarded as almost specific."

The waters in cases of six gallons, five dollars per case at the Springs. Springs open first day of June. Springs pamphlet sent to any address. Both water and pamphlets may be had of JOHN WYETH & BRO., 1412 Walnut St., Philadelphia.

THOS. F. GOODE, Proprietor,

Buffalo Lithia Springs, Virginia.

A VALUABLE WORK OF REFERENCE FOR PHYSICIANS.

THOMAS'S COMPREHENSIVE Pronouncing Medical Dictionary.

CONTAINING THE

ETYMOLOGY AND SIGNIFICATION OF THE TERMS MADE USE OF IN
MEDICINE AND THE KINDRED SCIENCES.

WITH AN APPENDIX COMPRISING

A COMPLETE LIST OF ALL THE MORE IMPORTANT ARTICLES OF THE MATERIA MEDICA, AND EXPLANATION OF
THE LATIN TERMS AND PHRASES OCCURRING IN ANATOMY, PHARMACY, Etc.,

Together with the Necessary Directions for Writing Latin Prescriptions.

BY J. THOMAS, M.D., A.M.

Author of Lippincott's Dictionary of Biography and Mythology, etc.

Crown Svo. Extra Cloth, \$3.25. Sheep, \$3.75.

From D. HAYES AGNEW, M.D., Professor of Surgery in University of Pennsylvania, etc.

"I have examined Dr. Thomas's Medical Dictionary with no small degree of care. I have never seen more industry, research, and scholarship compressed into so small a compass. In my range of reading, I know of no book which combines so pre-eminently thoroughness with brevity, conciseness with perspicuity and copiousness without redundancy, as this work. I anticipate for it a success of no ordinary character."

From the AMERICAN DRUGGISTS' CIRCULAR, New York.

"Dr. Thomas's Medical Dictionary is both elementary and thorough; though containing many hints and explanations unnecessary to a thorough classical scholar, it exhibits a degree of learning and research which we are safe in saying is surpassed by no other work of the kind."

From the BUFFALO MEDICAL AND SURGICAL JOURNAL.

"The work is eminently suited to the wants of the general practitioner and medical student, and, considering its size and cost, is, no doubt, unsurpassed in value by any similar work. We especially recommend it to the favorable notice of all medical students. It will be found invaluable."

From the CINCINNATI LANCET.

"Dr. Thomas and the publishers have performed their work well in preparing this most capital Medical Dictionary for professional use."

* * * For sale by all Booksellers, or will be sent, postage paid, upon receipt of the price by

J. B. LIPPINCOTT & CO., Publishers.

715 and 717 MARKET ST., PHILADELPHIA, PA.

UNIVERSITY OF THE CITY OF NEW YORK.

MEDICAL DEPARTMENT.

410 East Twenty-Sixth St., opposite Bellevue Hospital, New York City.

THIRTY-EIGHTH SESSION.—1878-79.

FACULTY OF MEDICINE.

REV. HOWARD CROSBY, D.D., LL.D., Chancellor of the University.	WILLIAM H. THOMPSON, M.D., Professor of Materia Medica and Therapeutics.
ALFRED C. POST, M.D., LL.D., Professor Emeritus of Clinical Surgery; President of the Faculty.	J. W. S. ARNOLD, M.D., Professor of Physiology and Histology.
CHARLES INSLEE PARDEE, M.D., Professor of Diseases of the Ear; Dean of the Faculty.	JOHN T. DARBY, M.D., Professor of Surgery.
JOHN C. DRAPER, M.D., LL.D., Professor of Chemistry.	J. WILLISTON WRIGHT, M.D., Professor of Obstetrics and Diseases of Women and Children.
ALFRED L. LOOMIS, M.D., Professor of Pathology and Practice of Medicine.	FANEUIL D. WEISSE, M.D., Professor of Practical and Surgical Anatomy.
WILLIAM DARLING, A.M., M.D., F.R.C.S., Professor of Anatomy.	JOSEPH W. WINTER, M.D., Demonstrator of Anatomy.

POST-GRADUATE FACULTY.

D. B. ST. JOHN ROOSA, M.D., Professor of Ophthalmology.	HENRY G. PIFFARD, M.D., Professor of Dermatology.
WM. A. HAMMOND, M.D., Professor of Diseases of the Mind and Nervous System.	A. E. MACDONALD, M.D., Professor of Medical Jurisprudence.
STEPHEN SMITH, M.D., Professor of Orthopædic Surgery.	JOSEPH W. HOWE, M.D., Clinical Professor of Surgery.
J. W. S. GOULEY, M.D., Professor of Diseases of the Genito-Urinary System.	LEWIS A. STIMSON, M.D., Professor of Pathological Anatomy.
MONTROSE A. Pallen, M.D., Professor of Gynæcology.	

THE COLLEGIATE YEAR is divided into three sessions,—a Preliminary Session, a Regular Winter Session, and a Spring Session.

THE PRELIMINARY SESSION will commence September 18, 1878, and will continue until the opening of the Regular Winter Session. It will be conducted on the plan of that session.

THE REGULAR WINTER SESSION will commence on the second of October, 1878, and end about the first of March, 1879.

The location of the new College edifice being immediately opposite the gate of Bellevue Hospital, and a few steps from the ferry to Charity Hospital, Blackwell's Island, the Students of the University Medical College are enabled to enjoy the advantages afforded by these Hospitals with the least possible loss of time. The Professors of the practical Chairs are connected with the Hospitals, and the University Students are admitted to *all the Clinics* given therein, *free of charge*.

In addition to the daily Hospital Clinics, there are eight Clinics each week in the College Building. Five Didactic Lectures will be given daily in the College Building, and Evening Recitations will be conducted by the Professors of Chemistry, Practice, Anatomy, Materia Medica, etc., Physiology, Surgery, and Obstetrics, upon the subjects of their lectures.

THE SPRING SESSION embraces a period of twelve weeks, beginning in the first week of March and ending the last week of May. The daily Clinics, Recitations, and Special Practical Courses will be the same as in the Winter Session, and there will be Lectures on Special Subjects by the Members of the Post-Graduate Faculty.

THE DISSECTING-ROOM is open throughout the entire collegiate year; material is abundant, and it is furnished free of charge.

STUDENTS WHO HAVE STUDIED TWO YEARS, and who have attended two full courses of Lectures, may be admitted to examination in Chemistry, Anatomy, and Physiology, and, if successful, will be examined at the expiration of their full course of Study on Practice, Materia Medica and Therapeutics, Surgery, and Obstetrics; but those who prefer it may have all their examinations at the close of their full term.

FEES.

For Course of Lectures	\$140.00
Matriculation	5.00
Demonstrator's fee, including material for dissection	10.00
Graduation fee	30.00
Post-Graduate Certificate	30.00

For further particulars and circulars, address the Dean,

Prof. CHARLES INSLEE PARDEE, M.D.,

University Medical College, 410 East 26th St., New York City.

UNIVERSITY OF PENNSYLVANIA, MEDICAL DEPARTMENT.

113th ANNUAL SESSION, 1878-79.

The Higher plan of Medical Education is now enforced, requiring attendance upon three graded courses of five months each. Didactic instruction is increased. Bedside instruction of each student in Physical Diagnosis, Clinical Medicine and Surgery, and Gynaecology is provided. The clinical and laboratory facilities are unrivalled in America. Pharmacy, Analytical and Medical Chemistry, Histology, Normal and Morbid Anatomy, and Operative Surgery are taught practically.

A Dental Department has recently been instituted.

For particulars, see ANNOUNCEMENT, for which address

JAMES TYSON, M.D., Secretary,
P. O. Box 2838, Philadelphia, Pa.

LONG ISLAND COLLEGE HOSPITAL, BROOKLYN, N. Y.

Annual Announcement and Circular for 1877-78.

The READING AND RECITATION TERM will commence October 2, 1878, and close February 14, 1879.
The REGULAR TERM will open March 5, 1878, and close the last week in June following.

FACULTY OF THE COLLEGE.

SAMUEL G. ARMOR, M.D., LL.D., Professor of the Practice of Medicine, and Dean.	EDWARD SEAMAN BUNKER, M.D., Professor of Obstetrics.
GEORGE W. PLYMPTON, A.M., Professor of Chemistry.	CORYDON L. FORD, M.D., Professor of Anatomy.
JOSEPH H. RAYMOND, M.D., Professor of Physiology.	JARVIS S. WIGHT, M.D., Professor of Surgery, and Registrar.
DANIEL AVRES, M.D., LL.D., Emeritus Professor of Surgery.	JOHN D. RUSHMORE, M.D., Professor of Materia Medica.
ALEXANDER J. C. SKENE, M.D., Professor of Diseases of Women and Children.	

The Clinical advantages of the Long Island College Hospital are unsurpassed in this country.
For Circulars, address DEAN or REGISTRAR.

IMPORTANT TO EVERY PHYSICIAN!!

Vol. V. begins January 1, 1879; gentlemen subscribing for 1879 will receive the GAZETTE from the time of remitting their subscriptions to January 1, *Gratis*.

THE HOSPITAL GAZETTE.

A WEEKLY JOURNAL OF MEDICINE, SURGERY, AND THE COLLATERAL SCIENCES.

EDITED BY

EDWARD J. BERMINGHAM, A.M., M.D., and FREDERICK A. LYONS, A.M., M.D.
H. H. KANE, M.D., Associate Editor.

This journal is published regularly *every* Thursday in the chief medical centre of this continent, and each number presents a faithful *weekly* epitome of the progress of Medical Science. The following departments are represented: *Lectures*, EDITORIALS, *Original Articles*, HOSPITAL RECORDS, *Translations*, PERISCOPE, *Bibliography*, SOCIETY PROCEEDINGS, *Correspondence*, NEWS ITEMS AND NOTES.

Each number contains TWENTY LARGE CLOSELY PRINTED PAGES of the choicest and most valuable reading-matter, thus making *1040 pages annually*, which, owing to our very large circulation, we are enabled to furnish at the remarkably low subscription price of

\$2.00 PER YEAR, INCLUDING POSTAGE,

thus making it the cheapest journal published.

✂ We decline to notice requests for specimen copies, but shall be happy to give any gentlemen desiring it a trial subscription for three months for 50 cents, payable in advance. This plan will give those contemplating subscribing an opportunity to thoroughly examine the merits of the journal.

Address all communications to

THE HOSPITAL GAZETTE,
NEW YORK CITY.

BELLEVUE HOSPITAL MEDICAL COLLEGE,

CITY OF NEW YORK.

MEMBER OF THE AMERICAN MEDICAL COLLEGE ASSOCIATION.

SESSIONS OF 1878-79.

THE COLLEGIATE YEAR in this Institution embraces a preliminary Autumnal Term, the Regular Winter Session, and a Spring Session.

THE PRELIMINARY AUTUMNAL TERM for 1878-1879 will open on Wednesday, September 18, 1878, and continue until the opening of the Regular Session. During this term, instruction, consisting of didactic lectures upon special subjects and daily clinical lectures, will be given, as heretofore, by the entire Faculty. Students expecting to attend the Regular Session are strongly recommended to attend the Preliminary Term, but attendance during the latter is not required. *During the Preliminary Term clinical and didactic lectures will be given in precisely the same number and order as in the Regular Session.*

THE REGULAR SESSION will begin on Wednesday, October 2, 1878, and end about the 1st of March, 1879.

FACULTY.

ISAAC E. TAYLOR, M.D.,

Emeritus Professor of Obstetrics and Diseases of Women, and President of the Faculty.

JAMES R. WOOD, M.D., LL.D.,

Emeritus Professor of Surgery.

FORDYCE BARKER, M.D.,

Professor of Clinical Midwifery and Diseases of Women.

AUSTIN FLINT, M.D.,

Professor of the Principles and Practice of Medicine and Clinical Medicine.

W. H. VAN BUREN, M.D.,

Professor of Principles and Practice of Surgery, Diseases of Genito-Urinary System, and Clinical Surgery.

LEWIS A. SAYRE, M.D.,

Professor of Orthopedic Surgery, and Clinical Surgery.

ALEXANDER E. MOTT, M.D.,

Professor of Clinical and Operative Surgery.

WILLIAM T. LUSK, M.D.,

Professor of Obstetrics and Diseases of Women and Children, and Clinical Midwifery.

WILLIAM M. POLK, M.D.,

Professor of Materia Medica and Therapeutics, and Clinical Medicine.

AUSTIN FLINT, JR., M.D.,

Professor of Physiology and Physiological Anatomy, and Secretary of the Faculty.

JOSEPH D. BRYANT, M.D.,

Professor of General, Descriptive, and Surgical Anatomy.

R. OGDEN DOREMUS, M.D., LL.D.,

Professor of Chemistry and Toxicology.

EDWARD G. JANEWAY, M.D.,

Professor of Pathological Anatomy and Histology, Diseases of the Nervous System, and Clinical Medicine.

PROFESSORS OF SPECIAL DEPARTMENTS, ETC.

HENRY D. NOYES, M.D.,

Professor of Ophthalmology and Otology.

JOHN P. GRAY, M.D., LL.D.,

Professor of Psychological Medicine and Medical Jurisprudence.

ERSKINE MASON, M.D.,

Clinical Professor of Surgery.

EDWARD L. KEYES, M.D.,

Professor of Dermatology, and Adjunct to the Chair of Principles of Surgery.

J. LEWIS SMITH, M.D.,

Clinical Professor of Diseases of Children.

LEROY MILTON YALE, M.D.,

Lecturer Adjunct upon Orthopedic Surgery.

A distinctive feature of the method of instruction in this College is the union of clinical and didactic teaching. All the lectures are given within the Hospital grounds. During the Regular Winter Session, in addition to four didactic lectures on every weekday except Saturday, two or three hours are daily allotted to clinical instruction.

The Spring Session consists chiefly of Recitations from Text-books. This term continues from the first of March until the first of June. During the Session daily recitations in all the departments are held by a corps of examiners appointed by the Faculty. Regular clinics are also given in the Hospital and in the College building.

FEES FOR THE REGULAR SESSION.


Fees for Tickets to all the Lectures during the Preliminary and Regular Term, including Clinical Lectures,	\$140.00
Matriculation Fee	5.00
Demonstrator's Ticket (including material for dissection)	10.00
Graduation Fee	30.00

FEES FOR THE SPRING SESSION.

Matriculation (Ticket good for the following Winter)	\$5.00
Recitations, Clinics, and Lectures	35.00
Dissection (Ticket good for the following Winter)	10.00

Students who have attended two full Winter Courses of Lectures may be examined at the end of their second course upon Materia Medica, Physiology, Anatomy, and Chemistry, and, if successful, they will be examined at the end of their third course upon Practice of Medicine, Surgery, and Obstetrics only.

For the Annual Circular and Catalogue, giving regulations for graduation and other information, address Prof. AUSTIN FLINT, JR., Secretary, Bellevue Hospital Medical College.



J. H. GEMRIG,

MANUFACTURER OF

Surgical and Orthopædical Instruments, Trusses, etc.,

109 S. EIGHTH ST., below Chestnut, PHILADELPHIA.



Extract from a Report of Awards at the Centennial Exhibition, 1876.

"The Finest Assortment of Surgical Instruments. The Instruments manufactured by this Exhibitor are perfect in every respect; their exquisite finish, and perfect balance of the Instruments, place them at the head of all competitors."

ILLUSTRATED CATALOGUE SENT ON APPLICATION.

DELLUC'S BISCOTINE FOR INFANTS AND INVALIDS.

This excellent and healthy substance is prepared from a sweetened bread and other nutritious substances, reduced to a fine powder, so as to render them easily soluble in water or milk.

A similar preparation is known in France, under the name of *Biscotes*, Food for Infants and Invalids.

As an article of common diet for infants, particularly those suffering from delicate constitutions, weak intestines, or looseness of the bowels, it will be found to give health and strength with more certainty than the Crude substances now in use, and not, like them, liable to sour on the stomach.

DELLUC & CO., 635 Broadway, New York.

To preserve your Teeth, use only Delluc's Eau Angelique Tooth Wash. Sold Everywhere.

IN PRESS.

THE TREATMENT OF DIPHTHERIA,

BASED UPON A NEW ETIOLOGY AND PATHOLOGY.

By WILLIAM C. REITER, A.M., M.D.

12mo. CLOTH LIMP.

THE CLIMATE OF AMERICA.

Its Influence in Health and Disease. A Practical Guide for Physicians and Invalids.

INCLUDING AN ACCOUNT OF THE

MINERAL SPRINGS OF AMERICA.

By WILLIAM PEPPER, M.D.,

Professor of Clinical Medicine in the University of Pennsylvania, etc.

ASSISTED IN THE SECTION ON MEDICAL TOPOGRAPHY

By JOHN M. KEATING, M.D.,

Physician to the Philadelphia Hospital, etc.

J. B. LIPPINCOTT & CO., PUBLISHERS.

NEW AND STANDARD MEDICAL BOOKS,

PUBLISHED BY

J. B. LIPPINCOTT & CO., PHILADELPHIA.

SCHWEIGGER'S HAND-BOOK OF OPHTHALMOLOGY.

By Prof. C. SCHWEIGGER, of the University of Berlin. Translated from the Third German Edition, by PORTER FARLEY, M.D. With Diagrams and other Illustrations. Svo. Extra cloth. \$4.50.

SAYRE'S SPINAL DISEASE AND SPINAL CURVATURE.

Their Treatment by Suspension, and the Use of the Plaster-of-Paris Bandage. By LEWIS A. SAYRE, M.D., Professor of Orthopaedic Surgery in Bellevue Hospital Medical College, etc. Handsomely Illustrated with Twenty-one Photographs from Nature, and numerous Woodcuts. 12mo. Extra cloth. \$4.00.

MITCHELL'S FAT AND BLOOD,

And How to Make Them. By S. WEIR MITCHELL, M.D., author of "Injuries of the Nerves," "Nurse and Patient," etc. 16mo. Fine cloth. \$1.25.

DUHRING'S DISEASES OF THE SKIN.

Diseases of the Skin. A Practical Treatise on Diseases of the Skin. By LOUIS A. DUHRING, M.D., author of "Atlas of Skin Diseases," Professor of Skin Diseases in the Hospital of the University of Pennsylvania, Physician to the Dispensary for Skin Diseases, Philadelphia, etc., etc. Svo. Illustrated. Extra cloth. \$6.00.

"We know of no work which the busy practitioner who has to treat this troublesome class of diseases can consult with more advantage, and we cordially recommend it."—*The British Medical Journal*.

"The most complete and satisfactory work on dermatology in the English language."—*Boston Medical and Surgical Journal*.

AITKEN ON THE SCIENCE AND PRACTICE OF MEDICINE.

Outlines of the Science and Practice of Medicine. By WILLIAM AITKEN, M.D. (Edin.), F.R.S., Professor of Pathology in the Army Medical School; Corresponding Member of the Imperial Society of Physicians of Vienna, etc. One vol. Svo. Cloth. \$5.00.

"It is well digested, clear, well written, and the work of a man who is conversant with every detail of his subject, and a thorough master of the art of teaching."—*British Medical Journal*.

ATLEE'S DIAGNOSIS OF OVARIAN TUMORS.

General and Differential Diagnosis of Ovarian Tumors, with Special Reference to the Operation of Ovariectomy. By WASHINGTON L. ATLEE, M.D. Illustrated. Svo. Extra cloth. \$5.00.

"In this work we have an unusual amount of *original* matter; in fact, it is altogether drawn from the author's long and carefully-studied experience. In this respect it is almost unique, and is an honor to independent American surgery."—*Philadelphia Medical and Surgical Reporter*.

DA COSTA'S MEDICAL DIAGNOSIS,

With Special Reference to Practical Medicine. A Guide to the Knowledge and Discrimination of Diseases. By J. M. DA COSTA, M.D., Lecturer on Clinical Medicine, and Physician to the Pennsylvania Hospital; Fellow of the College of Physicians of Philadelphia, etc., etc. *Fourth Edition, Revised*. With numerous Engravings. Svo. Cloth. \$6.00. Sheep. \$7.00.

"Da Costa's work is well known and highly and justly esteemed in England as in America. It is too firmly established, and its value too thoroughly recognized, to need a word *pro* or *con*."—*London Medical Times and Gazette*.

MITCHELL'S INJURIES OF THE NERVES,

And their Consequences. By S. WEIR MITCHELL, M.D. Cloth. \$3.00.


"Dr. Mitchell's well-earned reputation as an experimental physiologist of the highest order has been admirably maintained by the manner in which he has executed his task. . . . A volume which they can refer to now, and will be able to refer to in the future, with the utmost confidence and satisfaction."—*British and Foreign Medico-Chir. Rev.*

WOOD'S TREATISE ON THERAPEUTICS.

Comprising Materia Medica and Toxicology, with Especial Reference to the Application of the Physiological Action of Drugs to Clinical Medicine. By H. C. WOOD, Jr., M.D. *Second Edition, Revised and Enlarged*. Svo. Cloth. \$6.00. Sheep. \$6.50.

"Altogether, this work stands by itself as the only complete treatise on the physiological action of drugs in the English language, and no student of scientific therapeutics should be without it."—*London Practitioner*.

For sale by Booksellers generally, or will be sent by mail, postage paid, on receipt of the price by the Publishers.



GET THE STANDARD.

"It ought to be in every Library, also in every Academy and every School."—HON. CHARLES SUMNER.

WORCESTER'S QUARTO DICTIONARY.

A large, handsome volume of 1854 pages, containing considerably more than 100,000 Words in its Vocabulary, with the correct Pronunciation, Definition, and Etymology.

FULLY ILLUSTRATED AND UNABRIDGED. WITH FOUR FULL-PAGE ILLUMINATED PLATES. LIBRARY SHEEP, MARBLED EDGES, \$10.00.

"WORCESTER"

is now regarded as the standard authority, and is so recommended by BRYANT, LONGFELLOW, WHITTIER, SUMNER, HOLMES, IRVING, WINTHROP, AGASSIZ, MARSH, HENRY, EVERETT, MANN, QUINCY, FELTON, HILLARD, and the majority of our most distinguished scholars, and is, besides, recognized as authority by the Departments of our National Government.

THE COMPLETE SERIES OF WORCESTER'S DICTIONARIES.

Quarto Dictionary. Illustrated. Library Sheep. \$10.00.
Octavo (Universal and Critical) Dictionary. 8vo. Library Sheep. \$4.25.
Academic Dictionary. Crown 8vo. Half roan. \$2.00.
Comprehensive Dictionary. Illustrated. 12mo. Half roan. \$1.75.
School (Elementary) Dictionary. 12mo. Half roan. \$1.00.
Primary Dictionary. Illustrated. 16mo. Half roan. 60 cents.
Pocket Dictionary. Illustrated. 24mo. Cloth, 63 cents; Roan, flexible, 85 cents; Roan, tucks, gilt edges, \$1.00.

Many special aids to students, in addition to a very full pronouncing and defining vocabulary, make the above-named books, in the opinion of our most distinguished educators, the most complete as well as by far the cheapest Dictionaries of our language.

**For sale by all Booksellers, or will be sent, carriage free, on receipt of the price by

J. B. LIPPINCOTT & CO.,
PUBLISHERS, BOOKSELLERS, AND STATIONERS,
Nos. 715 and 717 MARKET STREET, PHILADELPHIA.

ALWAYS SPECIFY "K. & M." GRANULATED EFFERVESCENT SALTS.

ALWAYS SPECIFY "K. & M." GRANULATED EFFERVESCENT SALTS.

K. & M.
GRANULATED EFFERVESCENT
PEPSIN, BISMUTH, AND STRYCHNIA.
 ORIGINATED BY KEASBEY & MATTISON.

We again offer this preparation confidently to the profession. It has for four years enjoyed an extensive prescription among the most eminent of our practitioners. Its use has been attended with the most marked success perhaps of any of the various remedies recommended in dyspepsia, gastralgia, and kindred disorders. Its association with Carbonic Acid renders it palatable, and causes it to be retained by the most sensitive stomach. It is the only soluble form in which Pepsin and Bismuth should be ever given together, and is FIVE TIMES THE PEPSIN STRENGTH of the various elixirs, wines, liquors, etc., of the market. In reference to its reliability, we refer to the most prominent physicians in all sections of this country, who will very generally be found to be using it in their practice.

Always prescribe the GRANULATED EFFERVESCENT form, and always specify the "K. & M." Preparation.

Each drachm or heaping teaspoonful contains FIVE GRAINS Saccharated Pepsin, two grains soluble Citrate of Bismuth, and one-fiftieth grain Citrate of Strychnia.

Messrs. KEASBEY & MATTISON:

Gentlemen,—I have used a large number of your preparations for several years past with great satisfaction, and am happy to commend them to others from what I know both of your drugs and of yourselves.

Very truly yours,

W. W. KEEN, M.D.

K. & M.
GRANULATED EFFERVESCENT
CITRATE CAFFEINE.
 ORIGINATED BY KEASBEY & MATTISON.

The recent extensive employment of this salt as a remedy, amounting almost to a specific in cases of sick headache, and its use in nervous diseases generally, have caused us to place it upon our list. Its value is much enhanced by its administration with Carbonic Acid, so agreeably and efficiently produced by this form. *A heaping teaspoonful, containing ONE GRAIN of Citrate of Caffeine, should be given in sick headache, every hour or two before and during the paroxysms.*

Messrs. KEASBEY & MATTISON:

Gentlemen,—It gives me pleasure to add my testimony in regard to your preparations, which I have used for a long time; they are thoroughly reliable.

Truly yours,

PHILADELPHIA, September 27, 1877.

THOMAS G. MORTON, M.D.

K. & M.
GRANULATED EFFERVESCENT
CITRATE LITHIA.

In the effervescent form this chemical is administered in a peculiarly advantageous manner. Its association with Carbonic Acid renders it much less liable to disturb the stomach, while in strength it is greatly superior to Lithia Water. A valued diuretic and lithontriptic, its importance in the treatment of rheumatic, gouty, and analogous disorders can scarcely be over-estimated.

We have doubled the strength of this preparation, each drachm or heaping teaspoonful containing FOUR GRAINS of the chemically pure salt.

Messrs. KEASBEY & MATTISON:

Gentlemen,—I have employed your Extract of Malt and Infants' Food in a large number of cases, much to my own satisfaction and to the benefit of the patient. The Infants' Food I regard as especially useful in cases of infants requiring artificial food from any cause.

PHILADELPHIA, September 25, 1877.

Yours, etc.,

W. B. ATKINSON, M.D., Secretary American Medical Association.

K. & M.
GRANULATED EFFERVESCENT PREPARATIONS

of Magnesium Citrate, Potassium Bicarb., Citrate, Benzoate, and Bromide; Lithium Benzoate and Carbonate; Cerium Nitrate and Oxalate; Bismuth Citrate (SOLUBLE), Pepsin, and Bismuth; Pepsin, Bismuth, and Iron, etc., etc., etc.; also the following

K. & M.
GRANULATED EFFERVESCENT
MINERAL WATER SALTS.

Vichy, Kissingen, Seltzer, Carlshad, Congress, Crab Orchard, Pullna, Friedrichshall, etc., etc.

THE MANUFACTURERS,

KEASBEY & MATTISON,

Send for Pamphlet on New Remedies.

PHILADELPHIA.

A GREAT WORK ON SURGERY.

NEARLY READY.

The Science and Practice of Surgery.

SECOND EDITION. TWO VOLUMES.

1700 PAGES. 1000 ILLUSTRATIONS (MOSTLY ORIGINAL).

A Most Complete Text-Book for Practitioners and Students.

By FREDERICK JAMES GANT, F.R.S.,

Senior Surgeon and Professor of Surgery in the Royal Free Hospital,

ASSISTED BY THE FOLLOWING EMINENT AUTHORITIES:

Dr. BARNES, on the Surgery of the Female Genital Organs.

Prof. ERASMUS WILSON, F.R.S., on Diseases of the Skin.

Prof. HENRY POWER, F.R.S., on Ophthalmic Surgery.

Dr. LAIDLAW PURVES, on Aural Surgery.

Prof. TOMES, F.R.S., on Dental Surgery.

Dr. MORELL MACKENZIE, on Diseases and Injuries of the Throat.

Mr. W. ADAMS, F.R.C.S., on Orthopædic Surgery, etc., etc., etc.

The Publishers would invite the especial attention of the Profession to the above work. They claim it not only to be the latest, but one of the most complete books now to be had. The majority of the illustrations are original, and the essays of the various specialists on their particular branches add greatly to the interest of the volumes.

It is proposed to have it ready the latter part of September.

IN PRESS.

NAVAL HYGIENE. An Essay on the Various Influences Affecting Human Health, and of the Various Means of Preventing Disease, etc. By JOSEPH WILSON, M.D., Medical Director, U.S.N. Second Edition. Illustrated.

THE CELL DOCTRINE: Its History and Present State. By JAMES TYSON, M.D. Illustrated. Second Edition.

A PRACTICAL GUIDE TO THE EXAMINATION OF THE URINE. By JAMES TYSON, M.D. Illustrated. Second Edition.

LINDSAY & BLAKISTON,

MEDICAL PUBLISHERS,

25 South Sixth St., Philadelphia.

SYRUP-GLYCEROLE OF FERROUS CHLORIDE.

The Combination which all Forms of Iron must assume before Assimilation.

*Superior to all other Iron Salts in that, while it is readily taken into the circulation, and does re-ferri-
ginate the blood, it does not ordinarily cause constipation nor headache. It is
very pleasant to the taste, and is unalterable.*

Ferrous Chloride is a neutral salt, consequently it has no action upon the teeth, and may be admin-
istered fearlessly. This fact and the delightful taste of my preparation render it an advantageous form
for administration to children. Each fluidrachm contains one grain of Ferrous Chloride, which, owing
to the certainty of its assimilation, is a full, active, adult dose.

McKELWAY, Apothecary,

Successor to O. S. HUBBELL,

1410 CHESTNUT STREET.

VANITY FAIR

TOBACCO AND CIGARETTES.

TWO ARTICLES OF REAL MERIT.

Unsurpassed for PURITY, DELICACY, AND EXCELLENCE.

Their standard will ever be maintained.

Beware of Imitations and Counterfeits.

Sold Everywhere.

WM. S. KIMBALL & CO.,

Peerless Tobacco Works.

(5 First Prize Medals.)

Rochester, N. Y.

THE GREAT EUROPEAN NOVELTY.

HUNYADI JÁNOS.
NEW APERIENT WATER.



Specially recommended for
richness in aperient salts,
and its efficacy in Bilious
attacks, prevention of Gout,
Piles, etc., and as an ordi-
nary aperient, by LIEBIG,
VIRCHOW, SCANZO-
NI, and SIR HENRY
THOMPSON, and the
entire medical profession in
England and Germany.

APPROVED BY THE ACADEMY OF MEDICINE OF FRANCE, AND
ITS SALE IN FRANCE AUTHORIZED BY SPECIAL ORDER
OF THE FRENCH GOVERNMENT.

- DR. J. MARION SIMS, New York. "As a laxative,
I prefer it to every other mineral water."
DR. JAMES R. WOOD, New York. "Certain, but
gentle and painless; superior to any other bitter
water."
DR. WM. A. HAMMOND, New York. "The most
pleasant and efficient of all purgative waters."
DR. ALFRED L. LOOMIS, New York. "The most
prompt and most efficient; specially adapted for daily
use."
DR. FORDYCE BARKER, New York. "Requires
less, is less disagreeable and unpleasant than any
other."
DR. LEWIS A. SAYRE, New York. "Preferred to
any other laxative."

A WINEGLASSFUL A DOSE.

Indispensable to the travelling public.

Every genuine bottle bears the name of THE APOLLINARIS
Co. (Limited), London.

FRED'K DE BARY & CO.,

41 & 43 Warren St., New York,

Sole Agents for United States and Canadas.

FOR SALE BY DEALERS, GROCERS, AND DRUGGISTS.
The Label on every genuine bottle is printed on BLUE paper.

APOLLINARIS
NATURAL

Mineral Water.

THE QUEEN OF TABLE WATERS.

HIGHLY EFFERVESCENT.

APPROVED BY THE ACADEMY OF MEDICINE OF FRANCE, AND
ITS SALE IN FRANCE AUTHORIZED BY SPECIAL ORDER
OF THE FRENCH GOVERNMENT.

- DR. LEWIS A. SAYRE. "A delightful beverage."
DR. WILLIAM A. HAMMOND. "Far superior to
Vichy, Seltzer, or any other."
DR. ALFRED L. LOOMIS. "Most grateful and re-
freshing."
DR. R. OGDEN DOREMUS. "Absolutely pure and
wholesome; superior to all for daily use; free from all
the objections urged against Croton and artificially
aerated waters."
PROF. WANKLYN, London, Eng. "Impregnated
only with its own gas."
DR. E. R. PEASLEE. "Useful and very agreeable."
DR. AUSTIN FLINT. "Healthful, and well suited for
Dyspepsia, and cases of acute disease."
DR. JAMES R. WOOD. "Mildly antacid; agrees well
with dyspeptics, and where there is a gouty diathesis."
DR. FORDYCE BARKER. "By far the most agreeable;
alone or mixed with wine, useful in Catarrhs of Stomach
or Bladder and in Gout."
DR. J. MARION SIMS. "Not only a luxury, but a
necessity."

To be had of all Wine Merchants, Grocers, Drug-
gists, and Mineral Water Dealers through-
out the United States, and wholesale of

FRED'K DE BARY & CO.,

41 and 43 Warren Street,
NEW YORK.

Sole Agents for the United States.

Every genuine bottle bears the YELLOW Label.